

Inducing flocculation of non-floc-forming *Escherichia coli* cells

Yoshihiro Ojima, Masayuki Azuma, Masahito Taya

Citation	World Journal of Microbiology and Biotechnology, 34(12); 185
Issue Date	2018-12
Type	Journal Article
Textversion	author
Rights	This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: https://doi.org/10.1007/s11274-018-2563-z Springer Nature's AM terms of use: https://www.springernature.com/gp/open-research/policies/accepted-manuscript-terms
DOI	10.1007/s11274-018-2563-z

Self-Archiving by Author(s)
Placed on: Osaka City University

Mini-review: Inducing flocculation of non-floc-forming
Escherichia coli cells

Yoshihiro Ojima^{#1}, Masayuki Azuma¹, Masahito Taya²

¹*Department of Applied Chemistry and Bioengineering, Graduate School of Engineering, Osaka City University, 3-3-138, Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan*

²*Division of Chemical Engineering, Graduate School of Engineering Science, Osaka University, 1-3 Machikaneyama-cho, Toyonaka, Osaka 560-8531, Japan*

[#]Address correspondence to Yoshihiro Ojima, ojima@bioa.eng.osaka-cu.ac.jp.

Tel: +81-6-6605-2163

Keywords: *Escherichia coli*; microbial flocculation; non-floc-forming bacteria; flocculants

1 **Abstract**

2 The present article reviews several approaches for inducing flocculation of *Escherichia*
3 *coli* cells. The common industrially used bacterium *E. coli* does not naturally have
4 floc-forming ability. However, there are several approaches to induce flocculation of *E.*
5 *coli* cells. One is induction by flocculants— polyvalent inorganic salts, synthetic
6 polymeric flocculants, or bio-based polymeric materials, including polysaccharide
7 derivatives. Another method is the induction of spontaneous flocculation by changing
8 the phenotypes of *E. coli* cells; several studies have shown that physical treatment or
9 gene modification can endow *E. coli* cells with floc-forming ability. Coculturing *E. coli*
10 with other microbes is another approach to induce *E. coli* flocculation. These
11 approaches have particular advantages and disadvantages, and remain open to
12 clarification of the flocculation mechanisms and improvement of the induction
13 processes.

14 In this review, several approaches to the induction of *E. coli* flocculation are
15 summarized and discussed. This review will be a useful guide for the future
16 development of methods for the flocculation of non-floc-forming microorganisms.

1 **Introduction**

2 Flocculation is an aggregation phenomenon of microbial cells in which the cells form
3 flocs. The floc-forming capabilities of many microorganisms have been studied
4 ([Salehizadeh and Shojaosadati 2001](#)). In the 19th century, flocculation of the yeast
5 *Levure casseueese* was first reported by Louis Pasteur. Since then, flocculation in 19
6 other diverse microorganisms, including fungi and bacteria, has been confirmed
7 ([Nakamura et al. 1976](#)). In the activated sludge used in wastewater treatment and in
8 pure laboratory cultures, the components of flocs typically include polysaccharides,
9 proteins, and polynucleotides. These bacterial flocs are susceptible to hydrolytic
10 enzymes, such as cellulases, proteases, and deoxyribonucleases ([Tago and Aida 1977](#)).
11 Although the complete mechanism for microbial flocculation remains unclear,
12 exopolymeric materials play a key role ([Salehizadeh and Shojaosadati 2001](#)).
13 Flocculation can also be applied in industrial fermentation. For example, a smart
14 process has been established for high-performance ethanol production from molasses
15 using flocculating *Saccharomyces cerevisiae* ([Morimura et al. 1997](#)). Using flocculating
16 yeast eliminates the costly centrifugation step required for cell recovery during repeated
17 batch ethanol fermentation.

18 However, not all microbial species can form flocs. For example, the commonly used
19 industrial bacterium *Escherichia coli* does not naturally have floc-forming ability.
20 Therefore, it is necessary to induce the flocculation of such non-floc-forming bacteria.
21 If the objective of flocculation is the removal or inactivation of *E. coli* cells from water,
22 forced flocculation using chemicals or synthetic polymers is effective and inexpensive.
23 If the *E. coli* flocs are to be used as biocatalysts, environmentally friendly or

1 spontaneous flocculation is desirable. Therefore, it is desirable to have as many choices
2 as possible of methods to flocculate *E. coli* cells for different purposes.

3 In this review, several approaches to induce flocculation of *E. coli* cells, a typical
4 non-floc-forming microorganism, are summarized and discussed.

5

6 **Approaches for flocculation of *E. coli* cells**

7 Figure 1 summarizes possible approaches for flocculation of non-floc-forming *E. coli*.
8 These methods can be divided into flocculation of *E. coli* only and flocculation of *E.*
9 *coli* with other microbes. There are two different approaches for flocculation of *E. coli*
10 only. The first is using flocculants, which is the standard approach to induce the
11 flocculation of non-floc-forming bacteria. Flocculants are categorized into three types—
12 inorganic chemicals, synthetic polymers, and bio-based polymers. The other approach is
13 by changing the phenotypes or properties of the *E. coli* cells to generate floc-forming
14 ability by means of physical treatment or gene modification. Changing the phenotypes
15 or properties is relatively new and is desirable for the biocatalytic application of *E. coli*
16 flocs. If using mixed microbial species, the coculture of *E. coli* cells with other
17 microbes that have a floc-forming ability is a strong tool for inducing flocculation. The
18 details of each approach are described below.

19

20 **Flocculants**

21 Flocculants are widely used in industrial processes, including wastewater treatment,
22 downstream processing, and food and fermentation processes. As shown in Table 1, the
23 flocculants used can be categorized into three groups; (i) polyvalent inorganic
24 flocculants such as aluminum sulfate; (ii) organic synthetic polymer flocculants, such as

1 polyethylene imine (PEI) and polyacrylamide derivatives; and (iii) bio-based polymer
2 flocculants, such as chitosan, starch derivatives, and other microbial flocculants
3 ([Salehizadeh and Shojaosadati 2001](#)).

4

5 *Inorganic chemicals*

6 Polyvalent inorganic cations react with water to form hydroxide complexes. These
7 complexes react with phosphorus and suspended particulates to form a relatively
8 insoluble mass, which settles because of many factors, including a reduction of
9 electrical charge. Bacterial cells are colloidal particulates and can be aggregated. In the
10 case of *E. coli* flocculation, polyvalent inorganic cations, such as aluminum nitrate
11 ([Rubin and Hanna 1968](#)) and aluminum sulfate ([Bulson et al. 1984](#)) have long been
12 known to be suitable flocculants (Table 1). Recently, compounds of rare earth elements,
13 including lanthanum chloride ([Zhang et al. 2010](#)) and cerium nitrate ([Chen et al. 2010](#))
14 has been reported to induce flocculation of *E. coli* cells with high efficiency. However,
15 the high cost of rare earth elements is a problem with this approach. Addition of
16 inorganic compounds is an attractive approach to remove *E. coli* cells from raw water
17 because of the easy handling and low cost.

18

19 *Synthetic polymers*

20 Organic synthetic polymers are the most typical flocculants used because of their high
21 flocculating efficiency and low cost. In particular, cationic polyelectrolytes can
22 drastically change the degree of flocculation at concentration of ppm. PEI is a typical
23 cationic polyelectrolyte and its mechanism of *E. coli* flocculation has been well studied
24 ([Treweek and Morgan 1977](#)). Based on adsorption experiments and electrophoretic

1 mobility and refiltration rate measurements, it was concluded that the primary
2 mechanism of flocculation was not polymer bridging, but adsorption coagulation. Small
3 doses of high molecular-weight PEI species contributed to the formation of a charge
4 mosaic on the oppositely charged *E. coli* cell surface and this resulted in producing
5 rapid flocculation. The adsorbed PEI molecules, not only neutralized the negative
6 surface charge at the adsorption sites, but also caused localized charge reversal because
7 of the presence of excess cationic segments. *E. coli* flocs induced by PEI have been
8 applied as biocatalysts by Zou et al. ([Zou et al. 2018](#)). In this study, recombinant *E. coli*
9 expressing *Acidovorax facilis* nitrilase was flocculated with PEI, followed by
10 cross-linking with glutaraldehyde to obtain cross-linked cell aggregates (CLCAs). The
11 CLCAs were investigated as biocatalysts in the regioselective biotransformation of
12 1-cyanocyclohexaneacetonitrile into 1-cyanocyclohexaneacetic acid. The results showed
13 that the half-life of the CLCAs was drastically extended compared with that of free
14 cells.

15 Flocculation of *E. coli* cells using other synthetic polymers has been summarized by
16 Barany et al. ([Barany and Szepesszentgyorgyi 2004](#)). In this work, nonionic and anionic
17 polymers, including polyethylene oxide, polyvinyl alcohol, carboxylchitin, neutral
18 polyacrylamide, hydrolyzed (anionic) polyacrylamide, and polyacrylic acid showed
19 weak flocculation of *E. coli* cells (less than 20% removal). In contrast, flexible cationic
20 polyelectrolytes, such as polydiethylaminoethylmethacrylate (polyDEAEMA) and
21 copolymers of polyDEAEMA with vinylpyrrolidone (polyDEAEMA/VP), acrylamide
22 (polyDEAEMA/AA), and acrylic acid (polyDEAEMA/AC), were excellent flocculants
23 of *E. coli* suspensions; the use of these polymers at concentrations of 15–20 $\mu\text{g}/10^9$ cells
24 precipitated 90% of *E. coli* cells. On the basis of complex measurements of polymer

1 adsorption and its effect on the electrokinetic potential and degree of aggregation of
2 cells, it was concluded that the aggregation of *E. coli* cells by polyDEAEMA and
3 copolymers was because of charge neutralization ([Barany and Szepesszentgyorgyi](#)
4 [2004](#)).

5

6 *Bio-based polymers*

7 Traditional flocculants, such as inorganic salts and synthetic polymers, have been
8 proven to provide high flocculation efficiencies in water without large amounts of
9 bacteria. Recently, bio-based flocculants, such as starch, cellulose, chitosan
10 ([Salehizadeh and Shojaosadati 2001](#)), and polyglutamic acid ([Liu et al. 2018](#); [Liu et al.](#)
11 [2017a](#)) have attracted increasing attention due to their environmentally friendly property,
12 biodegradability, and widespread availability. In particular, developing novel bio-based
13 polymer flocculants with multi-functionality is important. Traditional flocculants, such
14 as those that are inorganic metal-based as well as synthetic organic polymers, have no
15 evident sterilization activities. Moreover, they themselves carry health risks, because of
16 residual metal ions or the release of noxious polymeric monomers into the target water.
17 General bio-based polymer flocculants have been previously reviewed ([Salehizadeh and](#)
18 [Shojaosadati 2001](#); [Salehizadeh and Yan 2014](#)). Here, we focus on bio-based polymers
19 targeting flocculation of *E. coli* cells with known compositions (i.e., excluding
20 flocculants based on unclarified bio-based polymers or mixtures). Bio-based polymeric
21 flocculants for *E. coli* cells are summarized in Table 1.

22 Chitosan, the deacetylation product of chitin, appears to be one of the most promising
23 candidates. The unique properties of chitosan mainly arise from the primary amine
24 groups present on the macromolecular backbone. Under acidic conditions, the molecular

1 chains have a positive charge, suggesting that this biopolymer is quite efficient for
2 flocculating contaminants that have negative surface charges. This fundamental
3 property would clearly provide a benefit in bacterial removal because most bacteria,
4 including *E. coli*, normally carry negative charges on the outside of their cell walls
5 ([Agerkvist et al. 1990](#)). Furthermore, there has been unique research into the fractal
6 structures of flocs formed by chitosan in terms of fractal dimensions ([Tang et al. 2001](#)),
7 providing a measurement of how the bacteria in the flocs occupy space. Research has
8 also demonstrated that chitosan shows antibacterial activity. To increase the bactericidal
9 effect, quaternary ammonium salt-grafted carboxymethyl chitosan has been developed
10 for *E. coli* flocculation ([Yang et al. 2014](#)). This polymer has bactericidal action through
11 the breaking of bacterial cell walls by the grafted quaternary ammonium salts. Chitosan
12 has also been used for the flocculation of *E. coli* cells for biocatalytic applications.
13 Flocculation using chitosan within a wide range of molecular weights and degrees of
14 acetylation can achieve a useful immobilization. On the basis of this technique, *E. coli*
15 cells expressing an omega-transaminase were successfully reused in consecutive batch
16 reactions ([Rehn et al. 2013](#)). Despite a very high density of cells in the immobilized
17 preparation, and a fast reaction, diffusion limitation was minimal. Thus, the natural
18 polymer chitosan and its derivatives are highly effective, not only as tools for the
19 removal of bacterial cells from water, but also for the immobilization of bacterial cells
20 for biocatalytic applications.

21 However, the high cost of chitosan limits its practical applications in water treatment.
22 Starch is an abundant natural resource and much is cheaper than chitosan. Because
23 starch contains large numbers of hydroxyl groups on the saccharide rings, starch can be
24 easily modified chemically for use in various applications by the introduction of

1 different functional groups onto the backbone. Flocculation of *E. coli* cells has been
2 reported using carboxymethyl-starch-graft-aminomethylated-polyacrylamide ([Huang et](#)
3 [al. 2016](#)). Under suitable pH conditions, this flocculant both effectively removed
4 turbidity and disrupted *E. coli* cells. In recent work, cationized starch-based flocculants
5 (starch-3-chloro-2-hydroxypropyl triethyl ammonium chloride, St-CTA) containing
6 various quaternary ammonium salt groups on the starch backbone have been used for *E.*
7 *coli* flocculation ([Liu et al. 2017b](#)). St-CTA with a high degree of substitution of CTA
8 improved the removal of contaminants due to the strong cationic nature and the charge
9 naturalization flocculation effect. This flocculant showed better antibacterial effects on
10 *E. coli* cells than on *Staphylococcus aureus* cells, indicating that the thicker cell walls of
11 the Gram-positive bacterium *S. aureus* are harder to break than the walls of *E. coli* cells.
12 In addition, cationized starch-based flocculants substituted with
13 glycidyltrimethylammonium chloride was also reported ([El-Naggar et al. 2018](#)). This
14 flocculant has achieved the same flocculation efficiency of aluminum sulphate. As a
15 new bio-based polymer flocculant, lignin nanoparticles (L-NPs) assembled with gelatin
16 was proposed for the *E. coli* flocculation ([Yin et al. 2018](#)). Positive charge of gelatin is
17 the driving force for flocculation of L-NPs-gelatin complex.

18

19 **Change in cellular phenotypes or properties**

20 The addition of flocculants is the most common approach for the flocculation of *E. coli*
21 cells. However, there are other approaches to induce spontaneous flocculation, including
22 by changing the phenotype or properties of *E. coli*. Both physical and biological
23 approaches have been reported (Table 2). In the physical approach, an electric field was
24 applied to enteroaggregative *E. coli* (EAEC) cells. In the biological approach, gene

1 modification was effective in inducing flocculation of laboratory *E. coli* strains. Both
2 these approaches can maintain a higher survival rate of *E. coli* cells inside the floc
3 structure than that in flocs induced by flocculants. Therefore, spontaneous flocculation
4 induced by a phenotype change may be more suitable for application in fermentation
5 processes than using conventional flocculants.

6

7 *Physical approaches*

8 EAEC is a diarrheal pathogen defined by a characteristic aggregative adherence to host
9 cells. The EAEC042 strain is known to have important distinguishing properties, such
10 as the positively charged surface protein dispersin and aggregative adherence fimbria
11 ([Goochee et al. 1987](#); [Nataro et al. 1985](#); [Sheikh et al. 2002](#)). In normal cases,
12 fimbrial-mediated EAEC042 adhesion to surfaces leads to biofilm formation. However,
13 Kumar et al. have showed that application of transverse low magnitude alternating
14 current and direct current electric fields in a culture chamber stopped biofilm formation
15 on a glass substrate, and led to flocculation ([Kumar et al. 2011](#)). EAEC042 flocs
16 induced by an electric field were more than 200 µm in size with a heterogeneous
17 composition. Both the current and magnitude of the electric field were important
18 parameters for controlling the cell viability in those flocs. These findings show promise
19 for the use of electric fields, not only for the manipulation of bacterial flocs, but also for
20 the treatment of medical instruments in preventing aggregative adherence to surfaces.

21

22 *Biological approaches*

23 Ojima et al. have demonstrated self-generated flocculation of *E. coli* cells by
24 overexpressing the native *bcsB* gene, which encodes a component of transmembrane

1 cellulose synthase complexes ([Ojima et al. 2015](#)). The resulting flocs had a paper-like
2 structure that was stable. Various *E. coli* laboratory strains including K-12, B, and O
3 formed visible flocs (>1 mm) by overexpressing the *bcsB* gene. The presence of green
4 fluorescent protein (GFP)-expressing *E. coli* cells was confirmed within the floc
5 structure, suggesting that the *E. coli* cells inside the floc structure are likely to be alive.
6 The flocs were sensitive to proteinases, indicating that the main component linking the
7 flocs was proteinous. Both protein analyses and observations of the flocs by
8 transmission electron microscopy indicated the involvement of outer membrane vesicles
9 (OMVs) in the flocculation of *E. coli* cells. OMVs are extracellular vesicles produced
10 by Gram-negative bacteria and are spherical bilayered proteolipids with a diameter of
11 20–250 nm. OMVs contain outer membrane proteins and lipids, periplasmic proteins,
12 lipopolysaccharides, RNA, and DNA ([Lee et al. 2007](#)). Gram-negative bacterial
13 biofilms that have formed either *in vivo* or *in vitro* typically contain numerous OMVs
14 ([Mashburn-Warren et al. 2008](#)). It has been observed that the *degP*-deficient mutant
15 ($\Delta degP$) cells spontaneously flocculated without overexpression of *bcsB* ([Ojima et al.](#)
16 [2015](#)). The *degP* gene encodes a periplasmic protease and its deletion strongly enhances
17 OMV production in *E. coli* cells ([Schwechheimer and Kuehn 2013](#)). In contrast,
18 *bcsB*-induced *E. coli* flocculation was greatly suppressed by the deletion of the $\Delta dsbA$
19 or $\Delta dsbB$ gene; these mutants are known to have considerably decreased OMV
20 production ([Schwechheimer and Kuehn 2013](#)). These results demonstrate a correlation
21 between the spontaneous flocculation of *E. coli* and enhanced OMV production.
22 Compared with forced flocculation, self-generated *E. coli* flocs have an advantage for
23 application in fermentation processes, because the cells within the flocs maintain the
24 viability and activity required for use as biocatalysts. For example, ethanol-producing *E.*

1 *coli* KO11 cells were endowed with floc-forming ability by overexpression of the *bcsB*
2 gene, without adverse effects on ethanol production ([Ojima et al. 2016](#)). In this study,
3 the glucose concentration and culture temperature were important parameters for the
4 flocculation of ethanol-producing *E. coli*. Sedimentation tests showed that the *E. coli*
5 flocs completed sedimentation within 15 min after cessation of shaking, while
6 planktonic cells remained suspended. The advantages of using flocculating *E. coli*
7 KO11 in ethanol production were demonstrated in a repeated batch operation.

8 When examining the flocculation of *bcsB*-overexpressing *E. coli* cells, mass
9 spectrometry analyses indicated that the elongation factor Ts (Tsf) was dominant protein
10 among the floc proteins ([Ojima et al. 2015](#)). Tsf is known to promote the release of
11 guanosine diphosphate by forming an intermediate complex with another elongation
12 factor, Tu, which is involved in the elongation cycle of protein biosynthesis ([Zhang et al.](#)
13 [1997](#)). A fusion protein consisting of Tsf and GFP was shown to be expressed
14 throughout the whole floc structure, even in the spaces without *E. coli* cells ([Ojima et al.](#)
15 [2018](#)). The amount of Tsf-GFP reached approximately 15% (w/w) of the total floc
16 protein, suggesting that the design and synthesis of a fusion protein with Tsf would
17 enable the display of a recombinant target protein on the structure of an *E. coli* floc.
18 Displaying a recombinant protein on flocs is a promising technique to construct
19 artificial microbial flocs with desired functionalities.

20

21 **Mixed with other microbes (coculture)**

22 Table 3 shows reported coculture systems of *E. coli* with other microbes for inducing
23 flocculation. Flocculation of *S. cerevisiae* has been much investigated because of its
24 importance in the brewing industry ([Soares 2011](#)). Flocculation of *S. cerevisiae* results

1 from an interaction between a lectin-like protein and mannose residues located on the
2 yeast cell surface. The FLO1 gene, which encodes a cell wall protein, plays an
3 important role in yeast flocculation, which is inhibited by mannose but not by glucose.

4 Interestingly, Peng et al. found flocculation of *E. coli* cells when they investigated the
5 probiotic effect of yeast cells against diarrhea caused by pathogenic *E. coli* ([Peng et al.
6 1997](#)); *E. coli* cells flocculated in the supernatant of liquid cultures of an antidiarrheal *S.
7 cerevisiae* strain. This flocculation of *E. coli* cells was induced by a glycoprotein
8 released by the yeast cells.

9 Peng et al. also investigated the coflocculation of *E. coli* cells with a variety of yeast
10 cells ([Peng et al. 2001a](#)). The results showed that the *E. coli* strain JM109 strain
11 coflocculated with *Candida utilis* G3, *Dekkera bruxellensis* G1, *Hanseniaspora
12 guilliermondii* H60, *Kloeckera apiculata* K315, *S. cerevisiae* HG, and
13 *Schizosaccharomyces pombe* G21 strains, even though these yeasts are
14 non-floc-forming strains. In addition, the FLO1 deletion mutant of *S. cerevisiae* also
15 coflocculated with *E. coli* cells, suggesting that coflocculation of *E. coli* and yeast cells
16 is independent of any inherent floc-forming-ability of the yeast cells. *S. pombe* showed
17 much less coflocculation than the other yeasts. *S. pombe* is known to have
18 galactose-rich cell walls and the glycosylation mutant *gms1* Δ induced a remarkable
19 amount of coflocculation ([Peng et al. 2001b](#)). It was concluded that *E. coli* lectins may
20 have specificity for α -1-6- and α -1-3-linked mannose residues of *S. pombe*, but in
21 wild-type *S. pombe* these mannose residues are shielded by galactose residues.

22 Coaggregation of *E. coli* with other probiotic strains was also confirmed with
23 *Lactobacillus* spp. ([Ekmekci et al. 2009](#)). Coaggregation of *L. acidophilus* S1 with *E.
24 coli* ATCC11229 was observed under both aerobic and anaerobic conditions. The

1 coaggregation of the strains was greater at acidic pH and decreased after heat treatment.
2 Thus, the coflocculation or coaggregation abilities of probiotic bacteria might enable
3 them to form a barrier that prevents colonization of pathogenic bacteria on host cells.

4

5 **Potential applications of different flocculation approaches**

6 Table 4 summarizes the advantages and disadvantages of different approaches for
7 inducing *E. coli* flocculation in potential fundamental research and industrial
8 applications. Flocculation of *E. coli* cells using either inorganic chemicals or synthetic
9 polymers is relatively inexpensive and involves easy handling. These flocculants are
10 suitable for wastewater treatment. However, toxicity toward *E. coli* cells and
11 detrimental effects on human health are disadvantages for the application of these
12 flocculants in bioproduction, even though several biocatalytic reactions have been
13 proposed using synthetic polymer-induced *E. coli* flocs. Bio-based polymer flocculants
14 are biodegradable, which gives these flocculants environmentally friendly properties.
15 However, bio-based polymers are not the first choice for wastewater treatment because
16 the cost is relatively high compared with inorganic chemicals or synthetic polymers. A
17 promising application of bio-based polymers is for use in aquaculture feed. Taking
18 advantage of the biodegradability, bio-based polymers can be added to aquaculture
19 ponds to recycle non-utilized proteins and derivative microbial proteins including *E.*
20 *coli* (Avnimelech 2015). Flocculation of pathogenic *E. coli* cells by an electric field
21 inhibits the biofilm formation and does not need any added flocculants. However, this
22 method cannot be scaled up easily for bulk reactions. The electrical field method might
23 be applied for the treatment of medical instruments to prevent biofilm infection. Gene
24 modification induces flocculation, without cost or addition of flocculants, because the

1 growing cells spontaneously form flocs. However, this approach should only be applied
2 in closed reaction systems due to the legal limitations of handling of genetically
3 modified microorganisms. Repeated batch fermentation is one of the promising
4 applications using the sedimentation property of flocs. In addition, the engineering of
5 flocs by further gene modification is also effective in advancing biocatalytic reactions
6 from the laboratory to an industrial scale. Flocculation by coculture with *E. coli* has a
7 complex mechanism, which causes difficulty in the handling of the flocculation. In
8 fundamental research, the coculture approach might provide better understanding of
9 microbial interactions through flocculation. The coculture method is also applicable for
10 use in probiotics because the partner microbes remove the pathogenic *E. coli* cells from
11 the gut in the host animal by the flocculation.

12

13 **Conclusion**

14 Flocculation of non-floc forming bacteria using added flocculants has been
15 investigated and applied in wastewater treatment. In the case of *E. coli* flocculation,
16 inorganic-metal-based and synthetic organic polymeric flocculants are effective.
17 However, these traditional flocculants have no evident sterilization activities. Moreover,
18 they themselves carry health risks. Therefore, developing bio-based polymer flocculants
19 with multi-functionality is important. Several types of bio-based polymer flocculants
20 with antibacterial properties efficiently remove *E. coli* cells from water, although the
21 cost is still high. In contrast, flocculation by changing the phenotype of *E. coli* is a novel
22 and economical approach. In particular, such biological approaches can maintain a high
23 survival rate of *E. coli* cells inside the floc structure, and thus are suitable for
24 application in fermentation processes. In particular, flocculation induced by gene

1 modification is a promising technique to construct artificial microbial flocs with desired
2 functions. Coflocculation of *E. coli* and other microbes is a complex phenomenon;
3 coflocculation with probiotic bacteria might enable the formation of a barrier that
4 prevents colonization of pathogenic *E. coli* on host cells.

5

6 **Acknowledgments**

7 The authors acknowledge financial support from a Grant-in-Aid for Young Scientists
8 (B) (No. 17K14869) from the Ministry of Education, Culture, Sports, Science and
9 Technology of Japan, and a Chemical Innovation Encouragement Prize from the Japan
10 Association for Chemical Innovation. A draft of this manuscript was proofed by James
11 Allen, D. Phil, from Edanz Group (www.edanzediting.com/ac).

12

13 **Conflict of interest**

14 The authors declare that they have no conflict of interest.

15

16 **References**

17 Agerkvist I, Eriksson L, Enfors SO (1990) Selective flocculation with chitosan in

18 *Escherichia coli* disintegrates: effects of pH and nuclease treatment. Enzyme

19 Microb Technol 12(8):584-590

20 Avnimelech Y (2015) Biofloc Technology - A Practical Guidebook 3rd Edition. The

21 World Aquaculture Society, Baton Rouge, Louisiana, United States

22 Barany S, Szepesszentgyorgyi A (2004) Flocculation of cellular suspensions by

23 polyelectrolytes. Adv Colloid Interface Sci 111(1-2):117-129

24 Bulson PC, Johnstone DL, Gibbons HL, Funk WH (1984) Removal and inactivation of

1 bacteria during alum treatment of a lake. Appl Environ Microbiol 48(2):425-430

2 Chen A, Shi Q, Feng J, Ouyang Y, Chen Y, Tan S (2010) Dissociation of outer
3 membrane for *Escherichia coli* cell caused by cerium nitrate. J Rare Earth
4 28(2):312-315

5 Ekmekci H, Aslim B, Ozturk S (2009) Characterization of vaginal lactobacilli
6 coaggregation ability with *Escherichia coli*. Microbiol Immunol 53(2):59-65

7 El-Naggar ME, Samhan FA, Salama AAA, Hamdy RM, Ali GH (2018) Cationic starch:
8 Safe and economic harvesting flocculant for microalgal biomass and inhibiting *E.*
9 *coli* growth. Int J Biol Macromol 116:1296-1303

10 Goochee CF, Hatch RT, Cadman TW (1987) Some observations on the role of type 1
11 fimbriae in *Escherichia coli* autoflocculation. Biotechnol Bioeng 29(8):1024-1034

12 Huang M, Wang Y, Cai J, Bai J, Yang H, Li A (2016) Preparation of dual-function
13 starch-based flocculants for the simultaneous removal of turbidity and inhibition
14 of *Escherichia coli* in water. Water Res 98:128-137

15 Kumar A, Mortensen P, Mukherjee P, Retterer T, Dokytcz M (2011) Electric field
16 induced bacterial flocculation of enteroaggregative *Escherichia coli* 042. Appl
17 Phys Lett 98(25):253701

18 Lee EY, Bang JY, Park GW, Choi DS, Kang JS, Kim HJ, Park KS, Lee JO, Kim YK,
19 Kwon KH, Kim KP, Gho YS (2007) Global proteomic profiling of native outer
20 membrane vesicles derived from *Escherichia coli*. Proteomics 7(17):3143-53

21 Liu T, Nobeshima H, Ojima Y, Azuma M (2018) A new method to purify
22 poly- γ -glutamic acid using gemini quaternary ammonium salts and
23 characterization of its ionic complex. J Chem Eng Jpn 51(5):431-437

24 Liu T, Yamashita Y, Fukumoto T, Tachibana T, Azuma M (2017a) Flocculation of real

1 sewage sludge using polyglutamic acid produced by *Bacillus* sp. isolated from soil.
2 J Chem Eng Jpn 50(3):201-206

3 Liu Z, Huang M, Li A, Yang H (2017b) Flocculation and antimicrobial properties of a
4 cationized starch. Water Res 119:57-66

5 Mashburn-Warren L, McLean RJ, Whiteley M (2008) Gram-negative outer membrane
6 vesicles: beyond the cell surface. Geobiology 6(3):214-9

7 Morimura S, Ling ZY, Kida K (1997) Ethanol production by repeated-batch
8 fermentation at high temperature in a molasses medium containing a high
9 concentration of total sugar by a thermotolerant flocculating yeast with improved
10 salt-tolerance. J Ferment Bioeng 83(3):271-274

11 Nakamura J, Miyashiro S, Hirose Y (1976) Purification and chemical analysis of
12 microbial cell flocculant produced by *Aspergillus sojae* AJ 7002. Agr Biol Chem
13 40(3):619-624

14 Nataro JP, Baldini MM, Kaper JB, Black RE, Bravo N, Levine MM (1985) Detection of
15 an adherence factor of enteropathogenic *Escherichia coli* with a DNA Probe. J
16 Infect Dis 152(3):560-565

17 Ojima Y, Nguyen MH, Yajima R, Taya M (2015) Flocculation of *Escherichia coli* cells
18 in association with enhanced production of outer membrane vesicles. Appl
19 Environ Microbiol 81(17):5900-5906

20 Ojima Y, Nunogami S, Azuma M, Taya M (2018) Displaying a recombinant protein on
21 flocs self-produced by *Escherichia coli* through fused expression with elongation
22 factor Ts. Enzyme Microbial Technol 108:21-25

23 Ojima Y, Takeda S, Taya M (2016) Floc formation of ethanol producing *Escherichia*
24 *coli* KO11 cells and its application to repeated batch operation. J Chem Eng Jpn


- 1 49(8):793-798
- 2 Peng X, Iserentant D, Verachtert H (1997) Effect of cultivation conditions on the
3 yeast-induced flocculation of *Escherichia coli*. Food Technol Biotechnol
4 35(4):243-247
- 5 Peng X, Sun J, Iserentant D, Michiels C, Verachtert H (2001a) Flocculation and
6 coflocculation of bacteria by yeasts. Appl Microbiol Biotechnol 55(6):777-781
- 7 Peng X, Sun J, Michiels C, Iserentant D, Verachtert H (2001b) Coflocculation of
8 *Escherichia coli* and *Schizosaccharomyces pombe*. Appl Microbiol Biotechnol
9 57(1-2):175-181
- 10 Rehn G, Grey C, Branneby C, Adlercreutz P (2013) Chitosan flocculation: an effective
11 method for immobilization of *E. coli* for biocatalytic processes. J Biotechnol
12 165(2):138-144
- 13 Rubin AJ, Hanna GP (1968) Coagulation of the bacterium *Escherichia coli* by
14 aluminum nitrate. Environ Sci Technol 2(5):358-362
- 15 Salehizadeh H, Shojaosadati SA (2001) Extracellular biopolymeric flocculants. Recent
16 trends and biotechnological importance. Biotechnol Adv 19(5):371-385
- 17 Salehizadeh H, Yan N (2014) Recent advances in extracellular biopolymer flocculants.
18 Biotechnol Adv 32(8):1506-1522
- 19 Schwechheimer C, Kuehn MJ (2013) Synthetic effect between envelope stress and lack
20 of outer membrane vesicle production in *Escherichia coli*. J Bacteriol
21 195(18):4161-4173
- 22 Sheikh J, Czczulin JR, Harrington S, Hicks S, Henderson IR, Le Bouguenec C,
23 Gounon P, Phillips A, Nataro JP (2002) A novel dispersin protein in
24 enteroaggregative *Escherichia coli*. J Clin Invest 110(9):1329-1337

- 1 Soares EV (2011) Flocculation in *Saccharomyces cerevisiae*: a review. J Appl Microbiol
2 110(1):1-18
- 3 Tago Y, Aida K (1977) Exocellular mucopolysaccharide closely related to bacterial floc
4 formation. Appl Environ Microbiol 34(3):308-314
- 5 Tang S, Ma Y, Sebastine IM (2001) The fractal nature of *Escherichia coli* biological
6 flocs. Colloids Surf B 20(3):211-218
- 7 Treweek GP, Morgan JJ (1977) The mechanism of *E. coli* aggregation by
8 polyethyleneimine. J Colloid Interface Sci 60(2):258-273
- 9 Yang Z, Degorce-Dumas JR, Yang H, Guibal E, Li A, Cheng R (2014) Flocculation of
10 *Escherichia coli* using a quaternary ammonium salt grafted carboxymethyl
11 chitosan flocculant. Environ Sci Technol 48(12):6867-6873
- 12 Yin H, Liu L, Wang X, Wang T, Zhou Y, Liu B, Shan Y, Wang L, Lü X (2018) A novel
13 flocculant prepared by lignin nanoparticles-gelatin complex from switchgrass for
14 the capture of *Staphylococcus aureus* and *Escherichia coli*. Colloids Surf A
15 Physicochem Eng Asp 545:51-59
- 16 Zhang Y, Riley LK, Lin M, Hu Z (2010) Lanthanum-based concentration and
17 microrespirometric detection of microbes in water. Water Res 44(11):3385-3392
- 18 Zhang Y, Sun V, Spremulli LL (1997) Role of domains in *Escherichia coli* and
19 mammalian mitochondrial elongation factor Ts in the interaction with elongation
20 factor Tu. J Biol Chem 272(35):21956-21963
- 21 Zou S-P, Huang J-W, Xue Y-P, Zheng Y-G (2018) Highly efficient production of
22 1-cyanocyclohexaneacetic acid by cross-linked cell aggregates (CLCAs) of
23 recombinant *E. coli* harboring nitrilase gene. Process Biochem 65:93-99
24

1

2 **FIGURE LEGEND**

3 Fig. 1 Categories of approaches for inducing flocculation of non-floc-forming *E. coli*
4 cells.



5

Table 1 Materials for flocculation of *E. coli*

Flocculants	Characteristics	References
Inorganic chemicals		
Aluminum sulfate	Reduction of electrical charge	Bulson et al. 1984
Aluminum nitrate	Reduction of electrical charge	Rubin and Hanna 1968
Lanthanum chloride	Reduction of electrical charge	Zhang et al. 2010
Cerium nitrate	Reduction of electrical charge	Chen et al. 2010
Synthetic polymers		
Polyethyleneimine (PEI)	Positive charge	Treweek and Morgan 1977
Polyethyleneimine (PEI)	Biocatalytic application	Zou et al. 2018
polydiethylaminoethylmetacrylate (polyDEAEMA) and its copolymers	Charge neutralization	Barany and Szepesszentgyorgyi 2004
Bio-based polymers		
Chitosan	Positive charge	Agerkvist et al. 1990
Chitosan	Characterization of fractal structure	Tang et al. 2001
Quaternary ammonium salt grafted carboxymethyl chitosan (CMC-g-PDMC)	Increased charge and bridging with bactericide	Yang et al. 2014
Chitosan	Biocatalytic application	Rehn et al. 2013
Carboxymethyl starch-grafted aminomethylated-polyacrylamide (CMS-g-APAM)	Positive charge with bactericide	Huang et al. 2016
Starch-3-chloro-2-hydroxypropyl triethyl ammonium chloride (St-CTA)	Positive charge with bactericide	Liu et al. 2017b
Cationized starch with glycidyltrimethylammonium chloride (GTAC)	Positive charge with bactericide	El-Naggar et al. 2018
Lignin nanoparticles assembled with gelatin	Positive charge	Yin et al. 2018

Table 2 Modification of cellular phenotypes or properties of *E. coli* for flocculation

Approaches	Mechanisms	References
Type1 fimbriae-positive cell at low pH	Fimbriae-to-fimbriae adhesion	Goochee et al. 1987
Electric field	Fimbriae-to-fimbriae adhesion	Kumar et al. 2011
Overexpression of <i>bcsB</i> gene	Proteinous component	Ojima et al. 2015
Deletion of <i>degP</i> gene	Proteinous component	Ojima et al. 2015

Table 3 Cocultures of *E. coli* with other microbes for flocculation

Strains	Phenomena	References
<i>S. cerevisiae</i> (antidiarrhea strain)	flocculation in the culture supernatant	Peng et al. 1997
<i>C. utilis</i> G3	coflocculation	Peng et al. 2001a
<i>D. bruxellensis</i> G1	coflocculation	Peng et al. 2001a
<i>H. guilliermondii</i> H60	coflocculation	Peng et al. 2001a
<i>K. apiculata</i> K315	coflocculation	Peng et al. 2001a
<i>S. cerevisiae</i> HG	coflocculation	Peng et al. 2001a
<i>S. pombe</i>	coflocculation	Peng et al. 2001b
<i>Lactobacillus</i> spp. (isolated from the lateral vaginal walls)	coaggregation	Ekmekci et al. 2009

Table 4 Advantages and disadvantages of different approaches for inducing *E. coli* flocculation in potential fundamental research and industrial applications

Approaches	Advantages	Disadvantages	Fundamental research	Industrial application
Inorganic chemicals	Low cost/ Easy handling	Toxicity	-	Wastewater treatment
Synthetic polymers	Easy handling	Toxicity	-	Wastewater treatment
Bio-based polymers	Environmentally friendly	High cost	-	Aquaculture feed/ Biocatalytic reactions
Electrical field	Easy handling/ Without flocculants	Small scale	-	Treatment of medical instruments
Gene modification	Low cost/ Without flocculants	Legal limitations of handling	Construction of engineered flocs	Repeated batch fermentation/ Biocatalytic reactions
Cocultures	Low cost/ Without flocculants	Difficult handling	Understanding of microbial interactions	Wastewater treatment/ Probiotics