Iridium-Catalyzed Direct Hydroarylation of Glycals via C–H Activation: Ligand-Controlled Stereoselective Synthesis of α - and β -C-Glycosyl Arenes

Kana Sakamoto, Masaki Nagai, Yusuke Ebe, Hideki Yorimitsu, Takahiro Nishimura

Citation	ACS Catalysis, 9(2); 1347-1352
Issue Date	2019-01-11
Type	Journal Article
Textversion	Author
Supporting Information	The Supporting Information is available free of charge on the ACS Publications website at https://doi.org/10.1021/acscatal.8b04686 . <ul style="list-style-type: none">• Experimental procedures and compound characterization (PDF)• X-ray data file (CIF)
Rights	This document is the Accepted Manuscript version of a Published Work that appeared in final form in ACS Catalysis, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see https://doi.org/10.1021/acscatal.8b04686 .
DOI	10.1021/acscatal.8b04686

Self-Archiving by Author(s)

Placed on: Osaka City University Repository

SAKAMOTO, K., NAGAI, M., EBE, Y., YORIMITSU, H., & NISHIMURA, T. (2019). Iridium-Catalyzed Direct Hydroarylation of Glycals via C–H Activation: Ligand-Controlled Stereoselective Synthesis of α - and β -C-Glycosyl Arenes. *ACS Catalysis*. 9, 1347-1352.

Iridium-Catalyzed Direct Hydroarylation of Glycals via C–H Activation: Ligand-Controlled Stereoselective Synthesis of α - and β -C-Glycosyl Arenes

Kana Sakamoto,[†] Masaki Nagai,[‡] Yusuke Ebe,[‡] Hideki Yorimitsu,[‡] and Takahiro Nishimura^{†,*}

[†] Department of Chemistry, Graduate School of Science, Osaka City University, Sumiyoshi, Osaka 558-8585, Japan

[‡] Department of Chemistry, Graduate School of Science, Kyoto University, Sakyo, Kyoto 606-8502, Japan

ABSTRACT: Diastereoselective direct hydroarylation of glycals with aromatic compounds was realized by use of an iridium/binap catalyst, giving the corresponding C-glycosyl compounds in high yields. The use of binap with the different absolute configuration enabled the stereoselective synthesis of α - or β -C-glycosyl arenes irrespective of the chirality of the glycals. The method showing the high functional group tolerance provides an atom-economical access to C-glycosyl arenes by way of C–H bond activation. **KEYWORDS:** iridium, hydroarylation, C–H activation, glycals, C-glycosyl arenes

C-Glycosyl derivatives have attracted much attention due to the existence of a number of natural products and the significant biological properties (Scheme 1).¹ For example, some C-glycosyl arenes as inhibitors of renal sodium-dependent glucose cotransporter 2 (SGLT2) have been recently approved to be drugs targeting type 2 diabetes.² In this context, many efforts have been devoted to provide efficient methods of introduction of aryl moieties into sugars for the synthesis of C-glycosyl arenes.¹ Transition metal-catalyzed cross-coupling reactions have played important roles in the C-glycosylations because the reactions often display regio- and/or stereoselectivity.³

Glycals represent sugar derivatives having a double bond between carbon atoms 1 and 2 of the ring, which allows installation of many functional groups into the monosaccharide. Pd-catalyzed Mizoroki–Heck type arylation of glycals has been extensively studied for the synthesis of unsaturated C-glycosyl compounds, where, in most cases, aryl groups were

introduced with α -selectivity.⁴ Ni-catalyzed reaction of arylboronic acids with perbenzylated glycals involving regioselective β -O elimination of the glycals was reported to give ring-opened products, which underwent the ring closure reactions leading to C-glycosyl arenes.⁵ Pd-catalyzed addition/elimination reactions of acylated glycals with uracils, which involves C–H activation, was recently reported.⁶ In this context, due to the growing medicinal importance of C-glycosyl arenes, a synthetic approach that uses the direct addition of arenes to glycals by C–H activation would be highly attractive from an atom-economical point of view.⁷

Recently, we reported Ir-catalyzed directed hydroarylation of vinyl ethers (Scheme 2a),⁸ where the aryl groups are selectively introduced into the α -carbon of the alkoxy group. The findings of the catalytic system of Ir prompted us to develop a new method for the synthesis of 2-deoxy aryl C-glycosyl derivatives in an atom-economical manner (Scheme 2b). Here we report the regio- and diastereoselective direct hydroarylation of glycals catalyzed by a chiral Ir complex.⁹ The stereoselective synthesis of C- α - or β -glycosyl arenes was achieved

Scheme 1. Selected Biologically Active C-Glycosyl Arenes

Scheme 2. Ir-Catalyzed Hydroarylation

Table 1. Ir/Binap-Catalyzed Stereoselective Hydroarylation of **2a with **1a**^a**

entry	ligand	temp. (°C)	yield (%) ^b	β:α ^c
1	(<i>S</i>)-binap	100	99	90:10
2	(<i>S</i>)-binap	80	90 ^d	96:4
3 ^e	(<i>S</i>)-binap	60	88	46:54
4	(<i>R</i>)-binap	100	84	12:88
5	(<i>R</i>)-binap	80	63	7:93
6 ^e	(<i>R</i>)-binap	60	83 ^d	5:95

^aReaction conditions: **1a** (0.11 mmol for entries 1–3, and 0.15 mmol for entries 4–6), **2a** (0.10 mmol), [IrCl(cod)]₂ (5 mol % Ir), binap (6 mol %), and NaBARF₄ (10 mol %) in toluene (0.40 mL) for 24 h. ^bCombined yield of the products determined by ¹H NMR using 1,4-dimethoxybenzene as an internal standard. ^cDetermined by ¹H NMR of the crude mixture. ^dIsolated yield of the major product. ^eFor 48 h.

by use of binap with the different absolute configuration.¹⁰

Initially, we selected glucal **2a** as an acceptor of the addition of a C–H bond of 2-phenylpyridine (**1a**) as shown in Table 1. After an optimization of the reaction conditions of the Ir catalysis being effective in our previous studies,^{8a} it was found that the diastereoselectivity in hydroarylation of glycals via C–H activation of aromatic compounds was controlled by changing the absolute configuration of a chiral ligand under the optimized reaction temperature. Thus, treatment of tri-*O*-acetyl-D-glucal (**2a**) with 1.1 equivalents of 2-phenylpyridine (**1a**) in the presence of [IrCl(cod)]₂ (5 mol % Ir, cod = 1,5-cyclooctadiene), (*S*)-binap¹¹ (6 mol %), and NaBARF₄ [Ar^F = 3,5-(CF₃)₂C₆H₃] (10 mol %) in toluene at 80 °C for 24 h gave C-glycosyl derivative **β-3aa** as a major product (β:α = 96:4, Table 1, entry 2).¹² Analytically pure **β-3aa** was isolated in 90% yield by preparative TLC. The high β-selectivity was observed at the high reaction temperature: the reaction at 60 °C gave an almost 1:1 mixture of the isomers (entry 3).¹³ The ligand-controlled β-selectivity of the present Ir catalysis is different from the selectivity observed in the reported Pd-catalysis of the addition/elimination reactions, which show α-selectivity significantly influenced by the stereochemistry of the glycals.⁴ In contrast, the use of (*R*)-binap enabled highly α-selective hydroarylation of D-glucal **2a** (entries 4–6), where the lower reaction temperature improved the α-selectivity. Thus, the reaction of **1a** (1.5 equiv.) with **2a** in the presence of the cationic Ir/(*R*)-binap catalyst at 60 °C for 48 h gave α-**3aa**

^aIsolated yield of the major product is shown unless otherwise noted. The diastereomeric ratio in the parenthesis was determined by ¹H NMR of the crude mixture. ^bAt 100 °C.

Scheme 3. Ir-Catalyzed β-Selective Hydroarylation of Glycals^a

as a major product (β:α = 5:95), which was isolated in 83% yield (entry 6).¹⁴

The protecting groups employed in the chemistry of sugars are highly important towards the preparation of complex oli-

^aIsolated yield of the major product is shown unless otherwise noted. The diastereomeric ratio in the parenthesis was determined by ¹H NMR of the crude mixture.

Scheme 4. Ir-Catalyzed α-Selective Hydroarylation of Glycals^a

Table 2. Ir/(*S*)-Binap-Catalyzed β -Selective Hydroarylation of **2f** with **1**^a

entry	1		yield (%) ^b	ratio (b:a) ^c
1		R = Me (1b)	86 (b- 3bf)	>99:1
2		MeO (1c)	83 (b- 3cf)	>99:1
3		F (1d)	72 (b- 3df)	>99:1
4		Br (1e)	73 (b- 3ef)	>99:1
5	Py = 2-pyridyl	Ph (1f)	87 (b- 3ff)	>99:1
6		R = OMe (1g)	66 (b- 3gf)	>99:1
7		Br (1h)	73 (b- 3hf)	>99:1
8		1i	77 (b- 3if)	>99:1
9		X = O (1j)	71 (b- 3jf)	>99:1
10 ^d		O (1j)	72 (b- 3jg)	>99:1
11		NMe (1k)	71 (b- 3kf)	>99:1

^aReaction conditions: **1** (0.15 mmol), **2f** (0.10 mmol), $[\text{IrCl}(\text{cod})]_2$ (5 mol % Ir), (*S*)-binap (6 mol %), and NaBARF_4 (10 mol %) in toluene (0.40 mL) at 80 °C for 24 h. ^bIsolated yield. ^cDetermined by ¹H NMR of the crude mixture. ^d**2g** (0.10 mmol) was used instead of **2f**.

gosaccharides. Therefore, a high functional group tolerance in the functionalization of sugars is desirable. Under the present catalytic conditions, diverse glycals **2b–h** participated in the β selective hydroarylation using (*S*)-binap to give high yields of the corresponding *C*-glycosyl arenes **3ab–ah** (Scheme 3). In the reaction of tri-*O*-benzyl-D-glucal (**2b**) and tri-*O*-benzyl-D-galactal (**2c**), a high β -selectivity was observed. Glucal **2d** having methoxymethyl groups reacted with **1a** to give **3ad** in 92% yield. Partially protected glucals **2e–g** can also be applied to the reaction to give the corresponding *C*-glycosyl arenes **3ae–ag** in high yields. Non-protected D-glucal (**2h**) also underwent the hydroarylation, but a slightly low diastereoselectivity was observed (β : α = 80:20).

Scheme 4 summarizes the results obtained for the α -selective hydroarylation of glycals using (*R*)-binap as a ligand. The hydroarylation of tri-*O*-benzyl-D-galactal (**2c**) and tri-*O*-acetyl-D-galactal (**2i**) took place with high α -selectivity. A good α -selectivity was also obtained in the reaction of mono-*O*-silylated **2e**. In contrast, α -selective reaction was not possible for **2f** giving β -**3af** as a major product probably due to the rigid conformation induced by a cyclic acetal moiety.¹⁵ The reaction of D-glucal (**2h**) also proceeded to give **3ah** in good yield and selectivity.

As shown in Schemes 3 and 4, the present catalytic system can be applied to a protecting group-free gulcal **2h**. However,

Table 3. Ir/(*R*)-Binap-Catalyzed α -Selective Hydroarylation of **2a** with **1**^a

entry	1		yield (%) ^b	ratio (b:a) ^c
1 ^d		R = Me (1b)	79 (a- 3ba)	4:96
2		MeO (1c)	91 (a- 3ca)	2:98
3		F (1d)	97 (a- 3da)	2:98
4	Py = 2-pyridyl	Br (1e)	84 (a- 3ea)	2:98
5		1h	67 (a- 3ha)	7:93
6 ^d		1i	67 (a- 3ia)	7:93
7		1j	65 (a- 3ja)	12:88
8		X = O (1j)	53 (a- 3ja)	1:99
9		S (1m)	56 (a- 3ma)	1:99

^aReaction conditions: **1** (0.15 mmol), **2a** (0.10 mmol), $[\text{IrCl}(\text{cod})]_2$ (5 mol % Ir), (*R*)-binap (6 mol %), and NaBARF_4 (10 mol %) in toluene (0.40 mL) at 60 °C for 48 h. ^bIsolated yield. ^cDetermined by ¹H NMR of the crude mixture. ^dFor 72 h.

the diastereoselectivity of the reaction is not enough high to obtain the pure isomers considering the difficulty of the purification. The problem was readily solved by deprotection of tri-*O*-acetyl groups in α - and β -**3aa** obtained in pure form. The deacetylation of α - and β -**3aa** in the presence of K_2CO_3 in methanol thus gave the corresponding unprotected *C*-glycosyl arenes α -**3ah** and β -**3ah** in 90% and 93% yields, respectively.

The present catalytic system can be applied to β -selective hydroarylation of glucal **2f** with several aromatic compounds **1** using (*S*)-binap (Table 2). 2-Phenylpyridines substituted with functional groups, Me (**1b**), MeO (**1c,g**), F (**1d**), Br (**1e,h**), and Ph (**1f**) are all good substrates to give the corresponding addition products **3bf–hf** with high diastereoselectivity (entries 1–7). An installation of a 3-thienyl group was also possible to give β -**3if** in 77% yield (entry 8). As directing groups, benzoxazolyl **1j** and benzimidazolyl **1k** functioned and the addition reactions of them gave the corresponding adducts in good yields (entries 9–11). Unfortunately, however, other aromatic compounds such as acetophenone, acetophenone *O*-methyl oxime, and *N*-(4-methoxyphenyl)-1-phenylethan-1-imine, which were good substrates for Ir-catalyzed *ortho*-C–H alkylation,^{8a,b} were inert under the present reaction conditions.

Scheme 5. Introduction of Two Sugar Units

α -Selective addition of a variety of aromatic groups to **2a** can be achieved using (*R*)-binap as the chiral ligand (Table 3). Aromatic compounds having a 2-pyridyl group (**1b–e,h,i**), 2-pyrimidyl (**1l**), 2-benzoxazolyl (**1j**), and benzothiazolyl (**1m**) were introduced into **2a** to give the corresponding α -C-glycosyl arenes in good yields (entries 1–9).

Finally, we conducted the installation of two sugar units into an aromatic molecule (Scheme 5). Thus, 1,3-di(2-pyridyl)benzene (**1n**) reacted with 2.2 equiv of glucal **2g** to give **4nf**, which involves two sugar units in the benzene ring (Scheme 5a). It should be noted that the *ortho*-C–H bond between two pyridyl rings, which is often metalated to give pin-cer metal complexes, is inert in the present reaction. The reaction of 4,4'-di(pyridin-2-yl)-1,1'-biphenyl (**1o**) was selectively alkylated by 2.2 equiv of glucal **2g** to give **4og** in 53% yield (Scheme 5b). The structure of **4og** was clarified by X-ray crystallographic analysis (Figure 1). These compounds are potential materials for multivalent glycoconjugates¹⁶ and further studies on the transformation of these compounds are underway.

In summary, we have developed the iridium-catalyzed regio- and diastereoselective hydroarylation of chiral glycals with aromatic compounds. The reaction proceeded via *ortho*-C–H activation of an aromatic ring and the subsequent regio- and stereoselective addition to glycals gave the hydroarylation products. The aryl groups were selectively introduced into the anomeric carbon and the stereoselectivity of the addition was controlled by the ligand. Thus, the use of binap with the different absolute configuration enabled the stereoselective synthesis of α - or β -C-glycosyl arenes irrespective of the chirality of the glycals. The present method provides an atom-economical access to C-glycosyl arenes via C–H bond activation.

ASSOCIATED CONTENT

AUTHOR INFORMATION

Corresponding Author

* Email: tnishi@sci.osaka-cu.ac.jp

Figure 1 ORTEP illustration of **4og** with thermal ellipsoids drawn at 30% probability level. Solvent molecules and hydrogen atoms are omitted for clarity. One of two independent molecules is shown.

Notes

The authors declare no competing financial interest.

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Experimental procedures and compound characterization (PDF and cif)

ACKNOWLEDGMENT

We acknowledge Prof. Loïc Lemiègre (Ecole Nationale Supérieure de Chimie de Rennes) for fruitful discussions. This work was supported by JSPS KAKENHI Grant No. 15H03810. Y.E. thanks the JSPS for a research Fellowship for Young Scientists. T.N. acknowledges Dr. Kenji Kitayama (Daicel Corporation) for fruitful discussions and Daicel Corporation for the financial support.

REFERENCES

- (1) For recent reviews, see: (a) Leng, W.-L.; Yao, H.; He, J.-X.; Liu, X.-W. Venturing beyond Donor-Controlled Glycosylation: New Perspectives toward Anomeric Selectivity. *Acc. Chem. Res.* **2018**, *51*, 628–639. (b) Kitamura, K.; Ando, Y.; Matsumoto, T.; Suzuki, K. Total Synthesis of Aryl C-Glycoside Natural Products: Strategies and Tactics. *Chem. Rev.* **2018**, *118*, 1495–1598. (c) Yang, Y.; Yu, B. Recent Advances in the Chemical Synthesis of C-Glycosides. *Chem. Rev.* **2017**, *117*, 12281–12356. (d) Bokor, É.; Kun, S.; Goyard, D.; Tóth, M.; Praly, J.-P.; Vidal, S.; Somsák, L. C-Glycopyranosyl Arenes and Hetarenes: Synthetic Methods and Bioactivity Focused on Antidiabetic Potential. *Chem. Rev.* **2017**, *117*, 1687–1764.
- (2) For selected examples, see: (a) Meng, W.; Ellsworth, B. A.; Nirschl, A. A.; McCann, P. J.; Petal, M.; Girotra, R. N.; Wu, G.; Sher, P. M.; Morrison, E. P.; Biller, S. A.; Zahler, R.; Deshpande, P. P.; Pullockaran, A.; Hagan, D. L.; Morgan, N.; Taylor, J. R.; Obermeier, M. T.; Humphreys, W. G.; Khanna, A.; Discenza, L.; Robertson, J. G.; Wang, A.; Han, S.; Wetterau, J. R.; Janovitz, E. B.; Flint, O. P.; Whaley, J. M.; Washburn, W. N. Discovery of Dapagliflozin: A Potent, Selective Renal Sodium-Dependent Glucose Cotransporter 2 (SGLT2) Inhibitor for the Treatment of Type 2 Diabetes. *J. Med. Chem.* **2008**, *51*, 1145–1149. (b) Nomura, S.; Sakamaki, S.; Hongu, M.; Kawanishi, E.; Koga, Y.; Sakamoto, T.; Yamamoto, Y.; Ueta, K.; Kimata, H.; Nakayama, K.; Tsuda-Tsukimoto, M. Discovery of Canagliflozin, a Novel C-Glucoside with Thiophene Ring, as Sodium-Dependent Glucose Cotransporter 2 Inhibitor for the Treatment of Type 2 Diabetes Mellitus. *J. Med. Chem.* **2010**, *53*, 6355–6360. (c) Imamura, M.; Nakanishi, K.; Suzuki, T.; Ikegai, K.; Shiraki, R.; Ogiyama, T.; Murakami, T.; Kurosaki, E.; Noda, A.; Kobayashi, Y.

- Yokota, M.; Koide, T.; Kosakai, K.; Ohkura, Y.; Takeuchi, M.; Tomiyama, H.; Ohta, M. Discovery of Ipragliflozin (ASP1941): A Novel C-Glycoside with Benzothiophene Structure as a Potent and Selective Sodium Glucose Co-transporter 2 (SGLT2) Inhibitor for the Treatment of Type 2 Diabetes Mellitus. *Bioorg. Med. Chem.* **2012**, *20*, 3263–3279. (d) Grempler, R.; Thomas, L.; Eckhardt, M.; Himmelsbach, F.; Sauer, A.; Sharp, D. E.; Bakker, R. A.; Mark, M.; Klein, T.; Eickelmann, P. Empagliflozin, A Novel Selective Sodium Glucose Cotransporter-2 (SGLT-2) Inhibitor: Characterisation and Comparison with Other SGLT-2 Inhibitors. *Diabetes Obes. Metab.* **2012**, *14*, 83–90. (e) Ohtake, Y.; Sato, T.; Kobayashi, T.; Nishimoto, M.; Taka, N.; Takano, K.; Yamamoto, K.; Ohmori, M.; Yamaguchi, M.; Takami, K.; Yeu, S.-Y.; Ahn, K.-H.; Matsuoka, H.; Morikawa, K.; Suzuki, M.; Hagita, H.; Ozawa, K.; Yamaguchi, K.; Kato, M.; Ikeda, S. Discovery of Tofogliflozin, a Novel C-Arylglycoside with an O-Spiroketal Ring System, as a Highly Selective Sodium Glucose Cotransporter 2 (SGLT2) Inhibitor for the Treatment of Type 2 Diabetes. *J. Med. Chem.* **2012**, *55*, 7828–7840. (f) Kakinuma, H.; Kakinuma, T.; Hashimoto-Tsuchiya, Y.; Arai, M.; Kawakita, Y.; Fukasawa, Y.; Iida, I.; Hagima, N.; Takeuchi, H.; Chino, Y.; Asami, J.; Okumura-Kitajima, L.; Io, F.; Yamamoto, D.; Miyata, N.; Takahashi, T.; Uchida, S.; Yamamoto, K. (1S)-1,5-Anhydro-1-[5-(4-ethoxybenzyl)-2-methoxy-4-methylphenyl]-1-thio-D-glucitol (TS-071) is a Potent, Selective Sodium-Dependent Glucose Cotransporter 2 (SGLT2) Inhibitor for Type 2 Diabetes Treatment. *J. Med. Chem.* **2010**, *53*, 3247–3261.
- (3) For reviews, see: (a) Li, X.; Zhu, J. Glycosylation via Transition-Metal Catalysis: Challenges and Opportunities. *Eur. J. Org. Chem.* **2016**, 4724–4767. (b) McKay, M. J.; Nguyen, H. M. Recent Advances in Transition Metal-Catalyzed Glycosylation. *ACS Catal.* **2012**, *2*, 1563–1595.
- (4) For a review, see: (a) Wellington, K. W.; Benner, S. A. A Review: Synthesis of Aryl C-Glycosides via the Heck Coupling reaction. *Nucleosides, Nucleotides, and Nucleic Acids* **2006**, *25*, 1309–1333. For selected examples, see: (b) Lei, M.; Gao, L.; Yang, J.-S. Microwave-Assisted Palladium-Catalyzed Cross-Coupling Reactions Between Pyranoid Glycals and Aryl Bromides. Synthesis of 2-Deoxy C-aryl- β -glycopyranosides, *Tetrahedron Lett.* **2009**, *50*, 5135–5138. (c) Li, H.-H.; Ye, X.-S. Regio- and Stereo-selective Synthesis of Aryl 2-Deoxy-C-glycopyranosides by Palladium-Catalyzed Heck Coupling Reactions of Glycals and Aryl Iodides, *Org. Biomol. Chem.* **2009**, *7*, 3855–3861. (d) Ramnauth, J.; Poulin, O.; Rakhit, S.; Madadaford, S. P. Palladium(II) Acetate Catalyzed Stereoselective C-Glycosidation of Peracetylated Glycals with Arylboronic Acids. *Org. Lett.* **2001**, *3*, 2013–2015. (e) Xiang, S.; Cai, S.; Zeng, J.; Liu, X.-W. Regio- and Stereoselective Synthesis of 2-Deoxy-C-aryl Glycosides via Palladium Catalyzed Decarboxylative Reactions. *Org. Lett.* **2011**, *13*, 4608–4611. (f) Bai, Y.; Li, Kim, M. H.; Liao, H.; Liug, X.-W. Oxidative Heck Reaction of Glycals and Aryl Hydrazines: A Palladium-Catalyzed C-Glycosylation. *J. Org. Chem.* **2013**, *78*, 8821–8825. (g) A. K. Kusunuru, S. K. Yousuf, M. Tatina, D. Mukherjee, Desulfative C-Arylation of Glycals by Using Benzenesulfonyl Chlorides. *Eur. J. Org. Chem.* **2015**, 459–462. (h) Tang, S.; Zheng, Q.; Xiong, D. C.; Jiang, S.; Li, Q.; Ye, X.-S. Stereoccontrolled Synthesis of 2-Deoxy-C-glycopyranosyl Arenes Using Glycals and Aromatic Amines. *Org. Lett.* **2018**, *20*, 3079–3082. For related examples of the Pd-catalyzed synthesis of C-glycosyl arenes, see: (i) Liu, M.; Niu, Y.; Wu, Y.-F.; Ye, X.-S. Ligand-Controlled Monoselective C-Aryl Glycoside Synthesis via Palladium-Catalyzed C–H Functionalization of *N*-Quinolyl Benzamides with 1-Iodoglycals. *Org. Lett.* **2016**, *18*, 1836–1839. (j) Zhang, S.; Niu, Y.-H.; Ye, X.-S. *Org. Lett.* **2017**, *19*, 3608–3611.
- (5) Liu, C.-F.; Xiong, D.-C.; Ye, X.-S. "Ring Opening–Ring Closure" Strategy for the Synthesis of Aryl-C-glycosides. *J. Org. Chem.* **2014**, *79*, 4676–4686.
- (6) F. Rasool; D. Mukherjee, Pd-Catalyzed Regio- and Stereoselective C-Nucleoside Synthesis from Unactivated Uracils and Pyranoid Glycals. *Org. Lett.* **2017**, *19*, 4936–4939.
- (7) For recent reviews on catalytic hydroarylation, see: (a) Dong, Z.; Ren, Z.; Thompson, S. J.; Xu, Y.; Dong, G. Transition-Metal-Catalyzed C–H Alkylation Using Alkenes. *Chem. Rev.* **2017**, *117*, 9333–9403. (b) Crisenza, G. E. M.; Bower, J. F. Branch Selective Murai-type Alkene Hydroarylation Reactions. *Chem. Lett.* **2016**, *45*, 2–9. (c) Yoshikai, N. Development of Cobalt-Catalyzed C–H Bond Functionalization Reactions. *Bull. Chem. Soc. Jpn.* **2014**, *87*, 843–857. (d) Zheng, Q.-Z.; Jiao, N. Transition-Metal-Catalyzed Ketone-Directed *Ortho*-C–H Functionalization Reactions. *Tetrahedron Lett.* **2014**, *55*, 1121–1126. (e) Rouquet, G.; Chatani, N. Catalytic Functionalization of C(sp²)–H and C(sp³)–H Bonds by Using Bidentate Directing Groups. *Angew. Chem., Int. Ed.* **2013**, *52*, 11726–11743. (f) Colby, D. A.; Tsai, A. S.; Bergman, R. G.; Ellman, J. A. Rhodium Catalyzed Chelation-Assisted C–H Bond Functionalization Reactions. *Acc. Chem. Res.* **2012**, *45*, 814–825. (g) Ackermann, L. Carboxylate-Assisted Transition-Metal-Catalyzed C–H Bond Functionalizations: Mechanism and Scope. *Chem. Rev.* **2011**, *111*, 1315–1345. (h) Colby, D. A.; Bergman, R. G.; Ellman, J. A. Rhodium-Catalyzed C–C Bond Formation via Heteroatom-Directed C–H Bond Activation. *Chem. Rev.* **2010**, *110*, 624–655.
- (8) (a) Ebe, Y.; Onoda, M.; Nishimura, T.; Yorimitsu, H. Iridium-Catalyzed Regio- and Enantioselective Hydroarylation of Alkenyl Ethers by Olefin Isomerization. *Angew. Chem., Int. Ed.* **2017**, *56*, 5607–5611. (b) Ebe, Y.; Nishimura, T. Iridium-Catalyzed Branch-Selective Hydroarylation of Vinyl Ethers via C–H Bond Activation. *J. Am. Chem. Soc.* **2015**, *137*, 5899–5902. (c) Hatano, M.; Ebe, Y.; Nishimura, T.; Yorimitsu, H. Asymmetric Alkylation of *N*-Sulfonylbenzamides with Vinyl Ethers via C–H Bond Activation Catalyzed by Hydroxo-iridium/Chiral Diene Complexes. *J. Am. Chem. Soc.* **2016**, *138*, 4010–4013. (d) Yamauchi, D.; Nishimura, T.; Yorimitsu, H. Asymmetric Hydroarylation of Vinyl Ethers Catalyzed by a Hydroxo-iridium Complex: Azoles as Effective Directing Groups. *Chem. Commun.* **2017**, 2760–2763.
- (9) For selected examples of Ir-catalyzed asymmetric hydroarylation, see: (a) Aufdenblatten, R.; Diezi, S.; Togni, A. ^[Ir]Iridium(I)-Catalyzed Asymmetric Intermolecular Hydroarylation of Norbornene with Benzamide. *Monatsh. Chem.* **2000**, *131*, 1345–1350. (b) Dorta, R.; Togni, A. Addition of the *Ortho*-C–H Bonds of Phenol Across an Olefin Catalyzed by a Chiral Iridium(I) Diphosphine Complex. *Chem. Commun.* **2003**, 760–761. (c) Tsuchikama, K.; Kasagawa, M.; Hashimoto, Y. ^[Ir]Endo, K.; Shibata, T. Cationic Iridium–BINAP Complex-Catalyzed Addition of Aryl Ketones to Alkynes and Alkenes via Directed C–H Bond Cleavage. *J. Organomet. Chem.* **2008**, *693*, 3939–3942. (d) Sevov, C. S.; Hartwig, J. F. Iridium-Catalyzed Intermolecular Asymmetric Hydroheteroarylation of Bicycloalkenes. *J. Am. Chem. Soc.* **2013**, *135*, 2116–2119. (e) Shirai, T.; Yamamoto, Y. Cationic Iridium/S-Me-BIPAM-Catalyzed Direct Asymmetric Intermolecular Hydroarylation of Bicycloalkenes. *Angew. Chem., Int. Ed.* **2015**, *54*, 9894–9897. (f) Shibata, T.; Michino, M. ^[Ir]Kurita, H.; Tahara, Y.; Kanyiva, K. S. Enantioselective Formal C–H Conjugate Addition of Acetanilides to β -Substituted Acrylates by Chiral Iridium Catalysts. *Chem. - Eur. J.* **2017**, *23*, 88–91. (g) Grélaud, S.; Cooper, P.; Feron, L. J.; Bower, J. F. Branch-Selective and Enantioselective Iridium-Catalyzed Alkene Hydroarylation via Anilide-Directed C–H Oxidative Addition. *J. Am. Chem. Soc.* **2018**, *140*, 9351–9356.
- (10) For reviews of catalyst-directed diastereoselective transformations, see: (a) Hoveyda, A. H.; Evans, D. A.; Fu, G. C. Substrate-Directable Chemical Reactions. *Chem. Rev.* **1993**, *93*, 1307–1370. (b) Mahatthananchai, J.; Dumas, A. M.; Bode, J. W. Catalytic Selective Synthesis. *Angew. Chem., Int. Ed.* **2012**, *51*, 10954–10990.
- (11) Takaya, H.; Mashima, K.; Koyano, K.; Yagi, M.; Kumobayashi, H.; Taketomi, T.; Akutagawa, S.; Noyori, R. Practical Synthesis of (*R*)- or (*S*)-2,2'-Bis(diarylphosphino)-1,1'-binaphthyls (BINAPs). *J. Org. Chem.* **1986**, *51*, 629–635.
- (12) The configurations were determined by ¹H NMR analysis. See the Supporting Information. The X-ray structure of **4og** also support the β -selectivity observed by using (*S*)-binap as a ligand.
- (13) The stereoselectivity of the addition is likely to be influenced by the conformation of the glycal, which may vary dependent on the reaction temperature.
- (14) Recent DFT calculations indicate that our Ir-catalyzed hydroarylation reactions of vinyl ethers proceed by carbometalation, see: (a) Zhang, M.; Huang, G. Mechanism of Iridium-Catalyzed Branched-Selective Hydroarylation of Vinyl Ethers: A Computational Study. *Dalton Trans.* **2016**, *45*, 3552. (b) Zhang, M.; Hu, L.; Lang, Y.; Cao,

Y.; Huang, G. Mechanism and Origins of Regio- and Enantioselectivities of Iridium-Catalyzed Hydroarylation of Alkenyl Ethers. *J. Org. Chem.* **2018**, *83*, 2937–2947. See also refs 8a and 8c for the mechanistic investigations.

(15) The rigid conformation induced by a cyclic acetal moiety may inhibit the formation of its ¹C₄ conformer of **α-3af**, which has the aryl group in an equatorial position. Rousseau, C.; Martin, O. R.

Stereodirected Synthesis of Aryl γ-C-Glycosides from 2-*O*-Arylsilyl-glucopyranosides. *Org. Lett.* **2003**, *5*, 3763–3766.

(16) (a) Cecioni, S.; Imbert, A.; Vidal, S. Glycomimetics versus Multivalent Glycoconjugates for the Design of High Affinity Lectin Ligands. *Chem. Rev.* **2015**, *115*, 525–561. (b) Chabre, Y. M.; Roy, R. Multivalent Glycoconjugate Syntheses and Applications Using Aromatic Scaffolds. *Chem. Soc. Rev.* **2013**, *42*, 4657–4708.