A SIMPLER EXPRESSION FOR COSTIN-MAZ'YA'S CONSTANT IN HARDY-LERAY INEQUALITY WITH WEIGHT

NAOKI HAMAMOTO

Citation	OCAMI Preprint Series. 20-9
Issue Date	2020-07-07
Type	Preprint
Textversion	Author
Rights	For personal use only. No other uses without permission.
Relation	This is a pre-print of an article published in Calculus of Variations and Partial Differential Equations. The final authenticated version is available online at: https://doi.org/10.1007/s00013-021-01595-z .

From: Osaka City University Advanced Mathematical Institute

<http://www.sci.osaka-cu.ac.jp/OCAMI/publication/preprint/preprint.html>

A SIMPLER EXPRESSION FOR COSTIN-MAZ'YA'S CONSTANT IN HARDY-LERAY INEQUALITY WITH WEIGHT

NAOKI HAMAMOTO

ABSTRACT. In this note, we obtain a simpler expression for the constant number given by Costin-Maz'ya [1] on sharp Hardy-Leray inequality for a class of solenoidal (namely divergence-free) vector fields, with respect to any radial power-weighted measure. The dependence of the constant on the weight exponent will be clear.

1. MOTIVATION AND MAIN RESULT

Throughout this paper, N is an integer with $N \geq 3$ and γ denotes any real number. What we call Costin-Maz'ya's constant is a real number expressed by the formula

$$C_{N,\gamma} = \left(\gamma + \frac{N}{2} - 1\right)^2 + \min \left\{ 2 + \min_{\tau \geq 0} \left(\tau + \frac{4(N-1)(\gamma-1)}{\tau + \left(\gamma - \frac{N}{2}\right)^2 + N-1} \right), N-1 \right\}. \quad (1.1)$$

It was found by Costin-Maz'ya [1] in the process of deriving the best constant in the weighted N -dimensional Hardy-Leray (or shortly “H-L”) inequality

$$C_{N,\gamma} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx \quad (1.2)$$

for solenoidal vector fields $\mathbf{u} = \mathbf{u}(\mathbf{x}) : \mathbb{R}^N \rightarrow \mathbb{R}^N$ (with a suitable regularity condition). This inequality serves as a solenoidal improvement of the original sharp H-L inequality

$$\left(\gamma + \frac{N}{2} - 1\right)^2 \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx$$

for unconstrained fields $\mathbf{u} : \mathbb{R}^N \rightarrow \mathbb{R}^N$, whose prototype case $\gamma = 0$ is famous for the one-dimensional inequality by Hardy [5] and its N -dimensional extension by Leray [6].

Practically, the derivation of the expression (1.1) by Costin-Maz'ya was carried out under the axisymmetry assumption on the solenoidal fields $\mathbf{u}$. Strictly speaking, however, their method includes incorrect datum for $N \geq 4$ overlooking the singular behavior of axisymmetric vector fields, and in fact the inequality (1.2) together with the expression (1.1) is invalid as long as the axisymmetry of $\mathbf{u}$ is assumed. (For details, see [2, §2.1].) Nevertheless, the validity of the same expression can be recovered if only we remove the axisymmetry assumption; indeed, in recent papers [4, 3] the author evaluated the constant $C_{N,\gamma}$ in the inequality (1.2) without any

2010 *Mathematics Subject Classification.* Primary 35A23; Secondary 26D10.

Key words and phrases. Hardy-Leray inequality, solenoidal fields, best constant, convex function, intermediate value theorem.

symmetry assumption on the solenoidal fields $\mathbf{u}$, and found its best value to be the same as in (1.1), or the lesser of the two real numbers

$$\begin{cases} C_{P,N,\gamma} := \left(\gamma + \frac{N}{2} - 1\right)^2 + 2 + \min_{\tau \geq 0} \left(\tau + \frac{4(N-1)(\gamma-1)}{\tau + \left(\gamma - \frac{N}{2}\right)^2 + N-1} \right), \\ C_{T,N,\gamma} := \left(\gamma + \frac{N}{2} - 1\right)^2 + N-1, \end{cases} \quad (1.3)$$

which are respectively the best constants in the H-L inequalities

$$\begin{cases} C_{P,N,\gamma} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx & \text{for poloidal fields } \mathbf{u}, \\ C_{T,N,\gamma} \int_{\mathbb{R}^N} \frac{|\mathbf{u}|^2}{|\mathbf{x}|^2} |\mathbf{x}|^{2\gamma} dx \leq \int_{\mathbb{R}^N} |\nabla \mathbf{u}|^2 |\mathbf{x}|^{2\gamma} dx & \text{for toroidal fields } \mathbf{u}, \end{cases}$$

in the sense of the so-called poloidal-toroidal decomposition of solenoidal fields.

In view of the observation above, Costin-Maz'ya's constant

$$C_{N,\gamma} = \min \{C_{P,N,\gamma}, C_{T,N,\gamma}\}$$

can be considered as meaningful, and hence would be better if its expression could be further simplified, although it has been regarded as unwieldy. Now, our goal is to get a simpler expression of $C_{N,\gamma}$ than (1.1); the statement of our main result reads as follows:

Theorem 1.1. *Let $C_{N,\gamma}$, $C_{P,N,\gamma}$ and $C_{T,N,\gamma}$ be the real numbers given in (1.1) and (1.3), and let $I_N = (\gamma_N^-, \gamma_N^+) \subset \mathbb{R}$ be the open interval between the two (extended) real numbers $\gamma_N^- < \gamma_N^+$ given by*

$$\gamma_N^- = \frac{N}{2} - \frac{N-1}{\sqrt{N+1}+2} \quad \text{and} \quad \gamma_N^+ = \begin{cases} \frac{N}{2} + \frac{N-1}{\sqrt{N+1}-2} & (N \geq 4) \\ \infty & (N = 3) \end{cases}.$$

Then it holds that

$$\begin{cases} C_{T,N,\gamma} < C_{P,N,\gamma} & \text{for } \gamma \in I_N, \\ C_{T,N,\gamma} = C_{P,N,\gamma} & \text{for } \gamma \in \{\gamma_N^-, \gamma_N^+\}, \\ C_{T,N,\gamma} > C_{P,N,\gamma} & \text{otherwise,} \end{cases}$$

and that $C_{P,N,\gamma} = \left(\gamma + \frac{N}{2} - 1\right)^2 \frac{\left(\gamma - \frac{N}{2}\right)^2 + N+1}{\left(\gamma - \frac{N}{2}\right)^2 + N-1}$ whenever $\gamma \notin I_N$. In particular,

$$\begin{aligned} C_{N,\gamma} &= \left(\gamma + \frac{N}{2} - 1\right)^2 + \min \left\{ 2 + \frac{4(N-1)(\gamma-1)}{\left(\gamma - \frac{N}{2}\right)^2 + N-1}, N-1 \right\} \\ &= \begin{cases} \left(\gamma + \frac{N}{2} - 1\right)^2 + N-1 & \text{for } \gamma \in I_N, \\ \left(\gamma + \frac{N}{2} - 1\right)^2 \frac{\left(\gamma - \frac{N}{2}\right)^2 + N+1}{\left(\gamma - \frac{N}{2}\right)^2 + N-1} & \text{otherwise.} \end{cases} \end{aligned}$$

It is clear that the expression of $C_{N,\gamma}$ in this theorem is simpler than (1.1). An advantage of such a simplification is that it helps us to verify that the equality sign in the inequality (1.2) is never achieved by any non-trivial solenoidal field $\mathbf{u}$. Indeed, in view of [3], one can observe that the main difficulty of this verification appears in the case where $C_{T,N,\gamma} > C_{P,N,\gamma}$ and simultaneously where the minimum

in (1.3) is not achieved by $\tau = 0$; the result of Theorem 1.1 tells us that such a case is void.

In the remaining of the present paper, we will prove Theorem 1.1, after preparing a technical lemma. The proof is elementary.

2. PROOF OF MAIN THEOREM

We start with proving the following fact, which plays a central role:

Lemma 2.1. *Let B be the quartic function given by*

$$B(\lambda) = \lambda^4 + (N-1)(2\lambda^2 - 4\lambda - (N-3)) \quad \forall \lambda \in \mathbb{R}.$$

Then it holds that $C_{T,N,\gamma} + 2 \leq C_{P,N,\gamma}$ whenever $B(\gamma - \frac{N}{2}) \leq 0$.

Proof. The key idea of the proof is an application of the intermediate value theorem to convex functions. First of all, notice that the convexity of B can be verified by the positivity of its second-order derivative:

$$B''(\lambda) = 12\lambda^2 + 4(N-1) > 0.$$

From this fact together with the datum

$$B(1 - \frac{N}{2}) = N^2/16 > 0 \quad \text{and} \quad B(1) = -(N-2)N < 0,$$

it follows that $\gamma > 1$ must be satisfied whenever $B(\gamma - \frac{N}{2}) \leq 0$.

Now, we set

$$F(\tau) := \tau + \frac{4(N-1)(\gamma-1)}{\tau + (\gamma - \frac{N}{2})^2 + N-1} - N + 3 \quad \forall \tau \geq 0 \quad (2.1)$$

in order that

$$\min_{\tau \geq 0} F(\tau) = C_{P,N,\gamma} - C_{T,N,\gamma}. \quad (2.2)$$

To evaluate the left-hand side, notice that a direct calculation yields

$$\frac{\partial F(\tau)}{\partial \tau} = \frac{\tau^2 + 2\left((\gamma - \frac{N}{2})^2 + N-1\right)\tau + B(\gamma - \frac{N}{2})}{\left(\tau + (\gamma - \frac{N}{2})^2 + N-1\right)^2}. \quad (2.3)$$

Then it follows from $B(\gamma - \frac{N}{2}) \leq 0$ that the minimum of F can be achieved by the nonnegative root of the equation $\partial F/\partial \tau = 0$ which we denote by

$$\begin{aligned} \tau_0 &:= -\left(\gamma - \frac{N}{2}\right)^2 - N + 1 + \sqrt{\left(\left(\gamma - \frac{N}{2}\right)^2 + N-1\right)^2 - B(\gamma - \frac{N}{2})} \\ &= -\left(\gamma - \frac{N}{2}\right)^2 - N + 1 + 2\sqrt{N-1}\sqrt{\gamma-1}. \end{aligned}$$

Therefore, the minimum value of $F - 2$ is given by the calculation

$$\min_{\tau \geq 0} F(\tau) - 2 = F(\tau_0) - 2 = -\left(\gamma - \frac{N}{2}\right)^2 - 2N + 2 + 4\sqrt{N-1}\sqrt{\gamma-1}.$$

Now, all that is left is to show the non-negativity of this value; it suffices to check the inequality

$$16(N-1)(\gamma-1) \geq \left(\left(\gamma - \frac{N}{2}\right)^2 + 2N-2\right)^2.$$

To this end, by putting $\lambda = \gamma - N/2$ we compute

$$\begin{aligned} & 16(N-1)(\gamma-1) - \left(\left(\gamma - \frac{N}{2}\right)^2 + 2N - 2\right)^2 + B\left(\gamma - \frac{N}{2}\right) \\ &= 16(N-1)\left(\lambda + \frac{N}{2} - 1\right) - \left(\lambda^2 + 2(N-1)\right)^2 + B(\lambda) \\ &= (N-1)(-2\lambda^2 + 12\lambda + 3N - 9). \end{aligned}$$

Hence, all we have to do is to check that the quadratic function

$$E(\lambda) := -2\lambda^2 + 12\lambda + 3N - 9$$

satisfies

$$E(\lambda) \geq 0 \quad \text{whenever} \quad B(\lambda) \leq 0.$$

For this purpose, we set

$$\lambda_{\pm} := 3 \pm \frac{\sqrt{6}}{2} \sqrt{N+3}$$

as the two roots of the quadratic equation $E(\lambda) = 0$. Then, with the aid of the polynomial division identity

$$B(\lambda) = -\frac{1}{4}(2\lambda^2 + 12\lambda + 7N + 59)E(\lambda) + 2(13N + 77)\lambda + \frac{1}{4}(N-3)(17N + 181),$$

we directly compute

$$\begin{cases} \lambda_- < 1 < \lambda_+, \\ B(1) = -(N-2)N < 0, \\ B(\lambda_+) = \frac{1}{4}(1305 + 442N + 17N^2) + \sqrt{6}\sqrt{N+3}(13N + 77) > 0, \\ B(\lambda_-) = \frac{1}{4} \frac{(N-3)^2(289N^2 + 538N - 503)}{17N^2 + 442N + 1305 + 4\sqrt{6}\sqrt{N+3}(13N + 77)} \geq 0. \end{cases}$$

In view of the concavity of E and the convexity of B , this fact tells us that

$$E(\lambda) \geq 0 \quad \text{holds whenever} \quad B(\lambda) \leq 0,$$

as desired. $\square$

Proof of Theorem 1.1. First of all, in view of (2.1), notice that the two numbers $\gamma_N^{\pm}$ are the roots of the equation in γ :

$$F(0) = \frac{4(N-1)(\gamma-1)}{\left(\gamma - \frac{N}{2}\right)^2 + N - 1} - N + 3 = 0.$$

Hence we see from (2.1) that

$$\begin{cases} F(0) > 0 & \text{for } \gamma \in I_N \\ F(0) = 0 & \text{for } \gamma \in \{\gamma_N^-, \gamma_N^+\}, \\ F(0) < 0 & \text{otherwise.} \end{cases} \quad (2.4)$$

Making use of Lemma 2.1 together with (2.2), we evaluate the sign of the number $C_{P,N,\gamma} - C_{T,N,\gamma}$ in the following cases:

The case $\gamma \in I_N$. When $B(\gamma - \frac{N}{2}) \leq 0$, we have $C_{T,N,\gamma} < C_{P,N,\gamma}$ by use of Lemma 2.1. When $B(\gamma - \frac{N}{2}) > 0$, the function $F(\tau)$ is monotone increasing in τ since its derivative (2.3) is positive for all $\tau \geq 0$; then the $\min_{\tau \geq 0} F(\tau)$ is attained at $\tau = 0$. Therefore, we see from (2.2) and (2.4) that

$$C_{P,N,\gamma} - C_{T,N,\gamma} = F(0) > 0,$$

and hence again that $C_{T,N,\gamma} < C_{P,N,\gamma}$.

The case $\gamma \in \mathbb{R} \setminus I_N$. We see from (2.2) that

$$C_{P,N,\gamma} - C_{T,N,\gamma} = \min_{\tau \geq 0} F(\tau) \leq F(0) \leq 0.$$

This fact together with Lemma 2.1 implies that $B(\gamma - \frac{N}{2})$ must be positive, whence we see in view of (2.3) that $F(\tau)$ is monotone increasing in $\tau \geq 0$. Therefore, we get $\min_{\tau \geq 0} F(\tau) = F(0)$ and

$$C_{P,N,\gamma} = C_{T,N,\gamma} + F(0) = \left(\gamma + \frac{N}{2} - 1\right)^2 \frac{\left(\gamma - \frac{N}{2}\right)^2 + N + 1}{\left(\gamma - \frac{N}{2}\right)^2 + N - 1}.$$

Moreover, recalling (2.4) we obtain
$$\begin{cases} C_{P,N,\gamma} = C_{T,N,\gamma} & \text{for } \gamma \in \{\gamma_N^-, \gamma_N^+\}, \\ C_{P,N,\gamma} < C_{T,N,\gamma} & \text{otherwise.} \end{cases}$$

The proof of Theorem 1.1 is now complete.

Acknowledgments

This research was partly supported by JSPS KAKENHI Grant-in-Aid for Scientific Research (B) 19H01800(F. Takahashi), and was also partly supported by Osaka City University Advanced Mathematical Institute (MEXT Joint Usage/Research Center on Mathematics and Theoretical Physics JPMXP0619217849).

REFERENCES

- [1] O. Costin and V. G. Maz'ya, *Sharp Hardy–Leray inequality for axisymmetric divergence-free fields*, Calculus of Variations and Partial Differential Equations **32** (2008), no. 4, 523–532.
- [2] N. Hamamoto, *Sharp Rellich–Leray inequality for axisymmetric divergence-free vector fields*, Calculus of Variations and Partial Differential Equations **58** (2019), no. 4, Paper No. 149, 23 pp.
- [3] ———, *Sharp Hardy–Leray inequality for solenoidal fields*, OCAMI Preprint Series (2020).
- [4] ———, *Three-dimensional sharp Hardy–Leray inequality for solenoidal fields*, Nonlinear Analysis **191** (2020), 111634, 14 pp.
- [5] G. H. Hardy, *Note on a theorem of Hilbert*, Mathematische Zeitschrift **6** (1920), no. 3–4, 314–317.
- [6] J. Leray, *Étude de diverses équations intégrales non linéaires et de quelques problèmes que pose l'hydrodynamique*, Journal de Mathématiques Pures et Appliquées **12** (1933), 1–82 (French).

OSAKA CITY UNIVERSITY ADVANCED MATHEMATICAL INSTITUTE
 3-3-138 SUGIMOTO, SUMIYOSHI-KU, OSAKA 558-8585, JAPAN
 E-mail address: yhjyoe@yahoo.co.jp (N. Hamamoto)