

BETA LAGUERRE ENSEMBLES IN GLOBAL REGIME

Hoang Dung Trinh and Khanh Duy Trinh

Citation	Osaka Journal of Mathematics. 58(2); 435-450
Issue Date	2021-04
Textversion	Publisher
Right	©Departments of Mathematics of Osaka University and Osaka City University.
DOI	10.18910/79433
Is Identical to	https://doi.org/10.18910/79433
Relation	The OJM has been digitized through Project Euclid platform http://projecteuclid.org/ojm starting from Vol. 1, No. 1.

Placed on : Osaka City University Repository

BETA LAGUERRE ENSEMBLES IN GLOBAL REGIME

HOANG DUNG TRINH and KHANH DUY TRINH

(Received September 20, 2019, revised January 16, 2020)

Abstract

Beta Laguerre ensembles, generalizations of Wishart and Laguerre ensembles, can be realized as eigenvalues of certain random tridiagonal matrices. Analogous to the Wishart ($\beta = 1$) case and the Laguerre ($\beta = 2$) case, for fixed β , it is known that the empirical distribution of the eigenvalues of the ensembles converges weakly to Marchenko–Pastur distributions, almost surely. The paper restudies the limiting behavior of the empirical distribution but in regimes where the parameter β is allowed to vary as a function of the matrix size N . We show that the above Marchenko–Pastur law holds as long as $\beta N \rightarrow \infty$. When $\beta N \rightarrow 2c \in (0, \infty)$, the limiting measure is related to associated Laguerre orthogonal polynomials. Gaussian fluctuations around the limit are also studied.

1. Introduction

Beta Laguerre (β -Laguerre) ensembles are ensembles of N positive particles distributed according to the following joint probability density function

$$(1) \quad \frac{1}{Z_{N,M}^{(\beta)}} \prod_{i < j} |\lambda_j - \lambda_i|^\beta \prod_{i=1}^N \left(\lambda_i^{\frac{\beta}{2}(M-N+1)-1} e^{-\frac{\beta M}{2} \lambda_i} \right), \quad (\lambda_i > 0),$$

where $\beta > 0$ and $M > N - 1$ are parameters, and $Z_{N,M}^{(\beta)}$ is the normalizing constant. For two special values $\beta = 1, 2$, they are the joint density of the eigenvalues of Wishart matrices and Laguerre matrices, respectively. For general $\beta > 0$, the ensembles can be realized as eigenvalues of a random tridiagonal matrix model $J_N = B_N(B_N)^t$ with a bidiagonal matrix B_N consisting of independent entries distributed as

$$B_N = \frac{1}{\sqrt{\beta M}} \begin{pmatrix} \chi_{\beta M} & & & & \\ \chi_{\beta(N-1)} & \chi_{\beta(M-1)} & & & \\ & & \ddots & \ddots & \\ & & & \ddots & \ddots \\ & & & & \chi_\beta & \chi_{\beta(M-N+1)} \end{pmatrix},$$

where χ_k denotes the chi distribution with k degrees of freedom, and $(B_N)^t$ denotes the transpose of B_N .

Wishart matrices (resp. Laguerre matrices) are random matrices of the form $M^{-1}G^tG$ (resp. $M^{-1}G^*G$), where G is an $M \times N$ matrix consisting of i.i.d. (independent identically distributed) entries of standard real (resp. complex) Gaussian distribution. Here G^* denotes the Hermitian conjugate of the matrix G . These random matrix models can be defined in a different way as invariant probability measures on the set of symmetric (resp. Hermitian)

matrices. The limiting behavior of their eigenvalues has been studied intensively, and it is known that as $N \rightarrow \infty$ with $N/M \rightarrow \gamma \in (0, 1)$, the empirical distribution

$$L_N = \frac{1}{N} \sum_{i=1}^N \delta_{\lambda_i}$$

converges to the Marchenko–Pastur distribution with parameter γ , almost surely. Here δ_λ denotes the Dirac measure at λ . The convergence means that for any bounded continuous function f ,

$$\int f(x) dL_N(x) = \frac{1}{N} \sum_{i=1}^N f(\lambda_i) \rightarrow \int f(x) mp_\gamma(x) dx \quad \text{as } N \rightarrow \infty, \text{ almost surely,}$$

where $mp_\gamma(x)$ is the density of the Marchenko–Pastur distribution with parameter γ ,

$$mp_\gamma(x) = \frac{1}{2\pi\gamma x} \sqrt{(\lambda_+ - x)(x - \lambda_-)}, \quad (\lambda_- < x < \lambda_+), \quad \lambda_\pm = (1 \pm \sqrt{\gamma})^2.$$

Gaussian fluctuations around the limit were also established with explicit formula for the limiting variance. The convergence to a limit and fluctuations around the limit of the empirical distributions are two typical problems in the global regime when a random matrix model is considered. Refer the book [12] for more details on Wishart and Laguerre matrices.

The convergence to Marchenko–Pastur distributions and fluctuations around the limit were extended to β -Laguerre ensembles for general $\beta > 0$ in [5] by using the random tridiagonal matrix model. Based on the model as well, results in the local regime and the edge regime were established [9, 14]. Note that the parameter β is fixed in those studies.

The aim of this paper is to refine results in the global regime of β -Laguerre ensembles. We assume that the parameter β varies as a function of N , while the parameter M depends on N in a way that $N/M \rightarrow \gamma \in (0, 1)$ as usual. We show that the Marchenko–Pastur law holds as long as $\beta N \rightarrow \infty$. When βN stays bounded, the limiting measure is related to associated Laguerre orthogonal polynomials. For the proof, we extend some ideas that were used in [7, 11, 15] in case of Gaussian beta ensembles. Our main results are stated in the following two theorems.

Theorem 1.1 (Convergence to a limit distribution). (i) As $\beta N \rightarrow \infty$, the empirical distribution L_N converges weakly to the Marchenko–Pastur distribution with parameter γ , almost surely. Here $L_N = N^{-1} \sum_{i=1}^N \delta_{\lambda_i}$ is the empirical distribution of the β -Laguerre ensemble (1).

(ii) As $\beta N \rightarrow 2c \in (0, \infty)$, the sequence $\{L_N\}$ converges weakly to the probability measure $\nu_{\gamma,c}$, almost surely, where $\nu_{\gamma,c}$ is given in Theorem 3.3.

To prove the above results, it is sufficient to show that any moment of L_N converges almost surely to the corresponding moment of $\nu_{\gamma,c}$ because the limiting measure $\nu_{\gamma,c}$ is determined by moments. Here for convenience, we refer to $\nu_{\gamma,\infty}$ as the Marchenko–Pastur distribution with parameter γ . It then follows that (see [6]) for any continuous function f of polynomial growth (there is a polynomial $P(x)$ such that $|f(x)| \leq P(x)$ for all $x \in \mathbb{R}$),

$$\int f(x) dL_N(x) = \frac{1}{N} \sum_{i=1}^N f(\lambda_i) \rightarrow \int f(x) d\nu_{\gamma,c}(x) \quad \text{as } N \rightarrow \infty, \text{ almost surely.}$$

Theorem 1.2 (Gaussian fluctuations around the limit). *Assume that the function f has continuous derivative of polynomial growth. Then the following hold.*

(i) As $\beta N \rightarrow \infty$,

$$\sqrt{\beta} \left(\sum_{i=1}^N f(\lambda_i) - \mathbb{E} \left[\sum_{i=1}^N f(\lambda_i) \right] \right) \xrightarrow{d} \mathcal{N}(0, \sigma_f^2),$$

where

$$\sigma_f^2 = \frac{1}{2\pi^2} \int_{\lambda_-}^{\lambda_+} \int_{\lambda_-}^{\lambda_+} \left(\frac{f(y) - f(x)}{y - x} \right)^2 \frac{4\gamma - (x - \lambda_m)(y - \lambda_m)}{\sqrt{4\gamma - (x - \lambda_m)^2} \sqrt{4\gamma - (y - \lambda_m)^2}} dx dy,$$

with $\lambda_m = (\lambda_- + \lambda_+)/2 = (1 + \gamma)$. Here ' \xrightarrow{d} ' denotes the convergence in distribution.

(ii) As $\beta N \rightarrow 2c \in (0, \infty)$,

$$\sqrt{\beta} \left(\sum_{i=1}^N f(\lambda_i) - \mathbb{E} \left[\sum_{i=1}^N f(\lambda_i) \right] \right) \xrightarrow{d} \mathcal{N}(0, \sigma_{f,c}^2),$$

where $\sigma_{f,c}^2$ is a constant.

The paper is organized as follows. In the next section, we introduce the random tridiagonal matrix model and related concepts. Results on convergence to a limit and Gaussian fluctuations around the limit are shown in Section 3 and Section 4, respectively.

2. Random tridiagonal matrix model and spectral measures

2.1. Random tridiagonal matrix model. Let G be an $M \times N$ random matrix consisting of i.i.d. real standard Gaussian random variables. Then $X = M^{-1}G'G$ is called a Wishart matrix. When $M \geq N$, the eigenvalues $\lambda_1, \dots, \lambda_N$ of X have the following joint density

$$\frac{1}{Z_{M,N}} |\Delta(\lambda)| \prod_{i=1}^N \left(\lambda_i^{\frac{1}{2}(M-N+1)-1} e^{-\frac{M}{2}\lambda_i} \right), \quad (\lambda_i > 0),$$

where $Z_{M,N}$ is the normalizing constant, and $\Delta(\lambda) = \prod_{i < j} (\lambda_j - \lambda_i)$ denotes the Vandermonde determinant. A Laguerre matrix $X = M^{-1}G^*G$ is obtained when the entries of G are i.i.d. of standard complex Gaussian distribution, where recall that G^* stands for the Hermitian conjugate of G . In this case, the joint density of the eigenvalues has an analogous formula to the Wishart case. Then β -Laguerre ensembles are defined to be ensembles of N positive particles with the joint density

$$(2) \quad \frac{1}{Z_{M,N}^{(\beta)}} |\Delta(\lambda)|^\beta \prod_{i=1}^N \left(\lambda_i^{\frac{\beta}{2}(M-N+1)-1} e^{-\frac{\beta M}{2}\lambda_i} \right), \quad (\lambda_i > 0),$$

where $\beta > 0$ and $M > N - 1$, which generalizes Wishart ($\beta = 1$) and Laguerre ($\beta = 2$) ensembles.

A random tridiagonal matrix model for β -Laguerre ensembles was introduced in [4] based on tridiagonalizing Wishart or Laguerre matrices. Let

$$B_N = \frac{1}{\sqrt{\beta M}} \begin{pmatrix} \chi_{\beta M} & & & & \\ \chi_{\beta(N-1)} & \chi_{\beta(M-1)} & & & \\ & & \ddots & \ddots & \\ & & & & \chi_{\beta} & \chi_{\beta(M-N+1)} \end{pmatrix}$$

denote a random bidiagonal matrix with independent entries. Then the eigenvalues of the tridiagonal matrix $J_N = B_N(B_N)^t$ are distributed as the β -Laguerre ensemble (2). Using this random matrix model and combinatorial arguments, the convergence to Marchenko–Pastur distributions and Gaussian fluctuations around the limit (only for polynomial test functions) were established in [5].

2.2. Spectral measures of Jacobi matrices. A symmetric tridiagonal matrix is called a Jacobi matrix. Spectral measures of Jacobi matrices $\{J_N\}$ have been studied recently. Here the spectral measure of J_N is defined to be the probability measure μ_N satisfying

$$\int x^k d\mu_N(x) = (J_N)^k(1, 1), \quad k = 0, 1, 2, \dots$$

It follows from the spectral decomposition of J_N that μ_N has the following form

$$\mu_N = \sum_{i=1}^N q_i^2 \delta_{\lambda_i}, \quad (q_i^2 = v_i(1)^2),$$

where v_1, \dots, v_N are the normalized eigenvectors of J_N corresponding to the eigenvalues $\lambda_1, \dots, \lambda_N$.

Spectral measures can also be defined for infinite Jacobi matrices. Let J be an infinite Jacobi matrix,

$$J = \begin{pmatrix} a_1 & b_1 & & & \\ b_1 & a_2 & b_2 & & \\ & & \ddots & \ddots & \ddots \end{pmatrix}, \quad a_i \in \mathbb{R}, b_i > 0.$$

Then there exists a probability measure μ such that

$$\int x^k d\mu(x) = J^k(1, 1), \quad k = 0, 1, \dots$$

When the above moment problem has a unique solution μ , or the measure μ is determined by moments, it is called the spectral measure of J . A simple, but useful sufficient condition for the uniqueness is given by [13, Corollary 3.8.9]

$$\sum_{n=1}^{\infty} \frac{1}{b_n} = \infty.$$

The matrix MP_{γ} in (3) and $J_{\alpha,c}$ in Lemma 2.1 clearly satisfy this condition. Thus, their spectral measures (the Marchenko–Pastur distribution with parameter γ and the measure $\mu_{\gamma,c}$ (and hence $\nu_{\gamma,c}$ in Theorem 3.3)) are all determined by moments.

By definition, moments of the spectral measure μ_N depend locally on upper left entries of J_N , and thus, the limiting behavior of μ_N follows easily from those of entries. In particular, for fixed β , as $N \rightarrow \infty$ with $N/M \rightarrow \gamma \in (0, 1)$, entry-wisely,

$$B_N = \frac{1}{\sqrt{\beta M}} \begin{pmatrix} \chi_{\beta M} & & & & \\ \chi_{\beta(N-1)} & \chi_{\beta(M-1)} & & & \\ & \ddots & \ddots & & \\ & & & \chi_{\beta} & \chi_{\beta(M-N+1)} \end{pmatrix} \rightarrow \begin{pmatrix} 1 & & & & \\ \sqrt{\gamma} & 1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \ddots \end{pmatrix}.$$

Here the convergence holds almost surely and in L_q for any $q \in [1, \infty)$. It follows that almost surely, the spectral measure μ_N converges weakly to the spectral measure of the following infinite Jacobi matrix

$$(3) \quad MP_{\gamma} = \begin{pmatrix} 1 & & & & \\ \sqrt{\gamma} & 1 & & & \\ & \ddots & \ddots & & \\ & & & \ddots & \ddots \end{pmatrix} \begin{pmatrix} 1 & \sqrt{\gamma} & & & \\ & 1 & \sqrt{\gamma} & & \\ & & \ddots & \ddots & \\ & & & \ddots & \ddots \end{pmatrix} = \begin{pmatrix} 1 & \sqrt{\gamma} & & & \\ \sqrt{\gamma} & 1 + \gamma & \sqrt{\gamma} & & \\ & \sqrt{\gamma} & 1 + \gamma & \sqrt{\gamma} & \\ & & \ddots & \ddots & \ddots \end{pmatrix},$$

which is nothing but the Marchenko–Pastur distribution with parameter γ [6].

For the Jacobi matrix J_N , the weights q_1^2, \dots, q_N^2 in the spectral measure μ_N are independent of the eigenvalues and have Dirichlet distribution with parameter $\beta/2$. From which the empirical distribution L_N and the spectral measure μ_N have the following relations

$$(4) \quad \mathbb{E}[\langle L_N, f \rangle] = \mathbb{E}[\langle \mu_N, f \rangle],$$

$$(5) \quad \text{Var}[\langle L_N, f \rangle] = \frac{\beta N + 2}{\beta N} \text{Var}[\langle \mu_N, f \rangle] - \frac{2}{\beta N} \left(\mathbb{E}[\langle \mu_N, f^2 \rangle] - \mathbb{E}[\langle \mu_N, f \rangle]^2 \right),$$

for suitable test functions f . Here we use the notation $\langle \mu, f \rangle$ to denote the integral $\int f d\mu$ for a measure μ and an integrable function f .

We conclude this section by giving several remarks on spectral measures of Jacobi matrices. The spectral measure μ orthogonalizes the sequence of polynomials $\{P_n\}_{n \geq 0}$ defined by

$$P_0 = 1, \quad P_1 = x - a_1, \\ P_{n+1} = xP_n - a_{n+1}P_n - b_n^2P_{n-1}, \quad n \geq 1.$$

In a particular case, $a_n = (\alpha + 2n - 1)$ and $b_n = \sqrt{n}\sqrt{\alpha + n}$, the sequence of polynomials $\{L_n\}$ defined by

$$L_0 = 1, \quad L_1 = x - (\alpha + 1), \\ L_{n+1} = xL_n - (\alpha + 2n + 1)L_n - n(\alpha + n)L_{n-1}, \quad n \geq 1$$

coincides with the sequence of scaled generalized Laguerre polynomials. The spectral measure in this case is the Gamma distribution with parameters $(\alpha + 1, 1)$, that is, the probability measure with density $\Gamma(\alpha + 1)^{-1}x^{\alpha}e^{-x}, x > 0$. In other words, the Gamma distribution with parameters $(\alpha + 1, 1)$ is the spectral measure of the following infinite Jacobi matrix J_{α} ,

$$J_{\alpha} = \begin{pmatrix} \alpha + 1 & \sqrt{\alpha + 1} & & & \\ \sqrt{\alpha + 1} & \alpha + 3 & \sqrt{2}\sqrt{\alpha + 2} & & \\ & & \ddots & \ddots & \ddots \end{pmatrix}$$

$$= \begin{pmatrix} \sqrt{\alpha+1} & & & \\ \sqrt{1} & \sqrt{\alpha+2} & & \\ & \ddots & \ddots & \\ & & \ddots & \ddots \end{pmatrix} \begin{pmatrix} \sqrt{\alpha+1} & \sqrt{1} & & \\ & \sqrt{\alpha+2} & \sqrt{2} & \\ & & \ddots & \ddots \end{pmatrix}.$$

When the entries of J_α are ‘shifted’ by a constant $c \in \mathbb{R}$, the resulting orthogonal polynomials are called associated Laguerre polynomials. The spectral measure in this case was explicitly calculated in [8] as Model II for associated Laguerre orthogonal polynomials.

Lemma 2.1 ([8]). *For $c > -1$ and $\alpha + c + 1 > 0$, let*

$$J_{\alpha,c} = \begin{pmatrix} \sqrt{\alpha+c+1} & & & \\ \sqrt{c+1} & \sqrt{\alpha+c+2} & & \\ & \ddots & \ddots & \\ & & \ddots & \ddots \end{pmatrix} \begin{pmatrix} \sqrt{\alpha+c+1} & \sqrt{c+1} & & \\ & \sqrt{\alpha+c+2} & \sqrt{c+2} & \\ & & \ddots & \ddots \end{pmatrix},$$

and $\mu_{\alpha,c}$ be its spectral measure. Then the density and the Stieltjes transform of $\mu_{\alpha,c}$ are given by

$$\begin{aligned} \mu_{\alpha,c}(x) &= \frac{1}{\Gamma(c+1)\Gamma(1+c+\alpha)} \frac{x^\alpha e^{-x}}{|\Psi(c, -\alpha; xe^{-i\pi})|^2}, \quad x \geq 0, \\ S_{\mu_{\alpha,c}}(z) &= \int_0^\infty \frac{\mu_{\alpha,c}(x)dx}{x-z} = \frac{\Psi(c+1, 1-\alpha; -z)}{\Psi(c, -\alpha; -z)}, \quad z \in \mathbb{C} \setminus \mathbb{R}. \end{aligned}$$

Here $\Psi(a, b; z)$ is Tricomi’s confluent hypergeometric function.

Note that when α is not an integer, an alternative formula for $\Psi(a, b; z)$ could be used

$$\begin{aligned} \Psi(c, -\alpha; xe^{-i\pi}) &= \frac{\Gamma(\alpha+1)}{\Gamma(\alpha+c+1)} {}_1F_1(c; -\alpha; -x) \\ &\quad - \frac{\Gamma(-\alpha-1)}{\Gamma(c)} x^{\alpha+1} e^{-i\pi\alpha} {}_1F_1(\alpha+c+1; 2+\alpha; -x), \end{aligned}$$

where ${}_1F_1(a; b; z)$ is the Kummer function.

3. Convergence to a limit distribution

In what follows, the parameter M depends on N in the way that $N/M \rightarrow \gamma \in (0, 1)$ as $N \rightarrow \infty$. We study the limiting behavior of the empirical distribution L_N through its moments $\langle L_N, x^r \rangle, r = 0, 1, 2, \dots$. Recall from the identity (4) that for $r = 0, 1, 2, \dots$,

$$\mathbb{E}[\langle L_N, x^r \rangle] = \mathbb{E}\left[\frac{1}{N} \text{tr}[(J_N)^r]\right] = \mathbb{E}[\langle \mu_N, x^r \rangle] = \mathbb{E}[(J_N)^r(1, 1)].$$

Here $\text{tr}[A]$ denotes the trace of a matrix A .

3.1. The Marchenko–Pastur regime, $\beta N \rightarrow \infty$. A key observation in this regime is the following asymptotic behavior of the chi distribution.

Lemma 3.1. *As $k \rightarrow \infty$,*

$$\frac{\chi_k}{\sqrt{k}} \rightarrow 1 \quad \text{in } L^q \text{ for any } 1 \leq q < \infty.$$

Let $\{c_i\}_{i=1}^N$ and $\{d_i\}_{i=1}^{N-1}$ be the diagonal and the sub-diagonal of B_N , respectively. Note that although we do not write explicitly, $\{c_i\}$ and $\{d_i\}$ depend on the triple (N, M, β) . It is clear

that as $\beta N \rightarrow \infty$,

$$B_N = \frac{1}{\sqrt{\beta M}} \begin{pmatrix} \chi_{\beta M} & & & & \\ \chi_{\beta(N-1)} & \chi_{\beta(M-1)} & & & \\ & \ddots & \ddots & & \\ & & & \chi_{\beta} & \chi_{\beta(M-N+1)} \end{pmatrix} \rightarrow \begin{pmatrix} 1 & & & & \\ \sqrt{\gamma} & 1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \ddots \end{pmatrix}.$$

Here the convergence means the pointwise convergence of entries, which holds in L^q for any $q \in [1, \infty)$, that is, for fixed i , as $N \rightarrow \infty$ with $\beta N \rightarrow \infty$,

(6)
$$c_i \rightarrow 1, \quad d_i \rightarrow \sqrt{\gamma} \quad \text{in } L^q \text{ for } q \in [1, \infty).$$

Since J_N is a tridiagonal matrix of the form

$$J_N = B_N(B_N)^t = \begin{pmatrix} c_1^2 & c_1 d_1 & & & \\ c_1 d_1 & c_2^2 + d_1^2 & c_2 d_2 & & \\ & \ddots & \ddots & \ddots & \\ & & & c_{N-1} d_{N-1} & c_N^2 + d_{N-1}^2 \end{pmatrix},$$

it follows that for fixed $r \in \mathbb{N}$, when $N > r$, $(J_N)^r(1, 1)$ is a polynomial of $\{c_i, d_i\}_{i \leq \frac{r+1}{2}}$. Let us show some explicit formulae for $(J_N)^r(1, 1)$,

$$\begin{aligned} J_N(1, 1) &= c_1^2, \\ (J_N)^2(1, 1) &= c_1^4 + c_1^2 d_1^2, \\ (J_N)^3(1, 1) &= c_1^6 + 2c_1^4 d_1^2 + c_1^2 c_2^2 d_1^2 + c_1^2 d_1^4. \end{aligned}$$

Then the pointwise convergence (6) implies that as $N \rightarrow \infty$ with $\beta N \rightarrow \infty$,

$$(J_N)^r(1, 1) \rightarrow (MP_\gamma)^r(1, 1) \quad \text{in } L^q \text{ for any } q \in [1, \infty),$$

where recall that the Jacobi matrix MP_γ is given in (3). Consequently, as $N \rightarrow \infty$,

$$\mathbb{E}[(J_N)^r(1, 1)] \rightarrow (MP_\gamma)^r(1, 1).$$

Recall also that the spectral measure of MP_γ is the Marchenko–Pastur distribution with parameter γ . Therefore, in this regime, the following convergence of the mean values holds.

Lemma 3.2. *For any $r \in \mathbb{N}$, as $N \rightarrow \infty$ with $\beta N \rightarrow \infty$,*

$$\mathbb{E}[\langle L_N, x^r \rangle] = \mathbb{E} \left[\frac{1}{N} \text{tr}[(J_N)^r] \right] = \mathbb{E}[(J_N)^r(1, 1)] \rightarrow \langle mp_\gamma, x^r \rangle.$$

3.2. High temperature regime, $\beta N \rightarrow 2c \in (0, \infty)$. Let

$$B_\infty := \begin{pmatrix} \tilde{\chi}_{\frac{2c}{\gamma}} & & & & \\ \tilde{\chi}_{2c} & \tilde{\chi}_{\frac{2c}{\gamma}} & & & \\ & \ddots & \ddots & & \\ & & & \ddots & \\ & & & & \ddots \end{pmatrix}, \quad \tilde{\chi}_k = \frac{1}{\sqrt{2}} \chi_k,$$

be an infinite bidiagonal matrix with independent entries. Then as $\beta N \rightarrow 2c$,

$$B_N = \frac{1}{\sqrt{\beta M}} \begin{pmatrix} \chi_{\beta M} & & & & & \\ \chi_{\beta(N-1)} & \chi_{\beta(M-1)} & & & & \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & \chi_{\beta} & \chi_{\beta(M-N+1)} \end{pmatrix} \xrightarrow{d} \frac{\sqrt{\gamma}}{\sqrt{c}} B_{\infty},$$

meaning that entries of B_N converge in distribution to the corresponding entries of the infinite matrix $\frac{\sqrt{\gamma}}{\sqrt{c}} B_{\infty}$. Recall that $(J_N)^r(1, 1)$ is a polynomial of $\{c_i, d_i\}_{i \leq \frac{r+1}{2}}$. Moreover, the entries of B_N and those of B_{∞} are independent. Therefore we can deduce that as $\beta N \rightarrow 2c$,

$$(J_N)^r(1, 1) \xrightarrow{d} (J_{\infty})^r(1, 1),$$

$$\mathbb{E}[(J_N)^r(1, 1)] \rightarrow \mathbb{E}[(J_{\infty})^r(1, 1)],$$

where

$$J_{\infty} = \frac{\gamma}{c} B_{\infty} (B_{\infty})^t.$$

By this approach, we have just shown the existence of the limit of the mean values when $\beta N \rightarrow 2c$. However, we are not able to identify the limit directly from J_{∞} .

To identify the limit, we now extend some ideas that used in [7] in case of Gaussian beta ensembles to show the following.

Theorem 3.3. *Let $\nu_{\gamma,c}$ be the spectral measure of the following Jacobi matrix*

$$\frac{\gamma}{c} \begin{pmatrix} \sqrt{\frac{c}{\gamma}} & & & & & \\ \sqrt{c+1} & \sqrt{\frac{c}{\gamma}+1} & & & & \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & \sqrt{c} & \sqrt{c+1} \\ & & & & \sqrt{\frac{c}{\gamma}+1} & \sqrt{c+2} \\ & & & & & \ddots & \ddots \end{pmatrix} = \frac{\gamma}{c} J_{\frac{\gamma}{\gamma}, c}.$$

Then for any $r \in \mathbb{N}$, as $N \rightarrow \infty$ with $\beta N \rightarrow 2c$,

$$\mathbb{E}[\langle L_N, x^r \rangle] = \mathbb{E} \left[\frac{1}{N} \text{tr}[(J_N)^r] \right] = \mathbb{E}[(J_N)^r(1, 1)] \rightarrow \langle \nu_{\gamma,c}, x^r \rangle.$$

Theorem 3.3 is equivalent to Theorem 3.5 below which states a result for scaled β -Laguerre ensembles. Let us switch to the scaled version. Let $\tilde{J}_N := \tilde{B}_N (\tilde{B}_N)^t$ be a Jacobi matrix, where

$$\tilde{B}_N = \begin{pmatrix} \tilde{\chi}_{2\alpha+2+2\kappa(N-1)} & & & & & \\ \tilde{\chi}_{2\kappa(N-1)} & \tilde{\chi}_{2\alpha+2+2\kappa(N-2)} & & & & \\ & & \ddots & & & \\ & & & \ddots & & \\ & & & & \tilde{\chi}_{2\kappa} & \tilde{\chi}_{2\alpha+2} \end{pmatrix},$$

(with $\kappa = \beta/2$ and $\alpha = \frac{\beta}{2}(M - N + 1) - 1 = \kappa(M - N + 1) - 1$). Then the joint density of the eigenvalues of \tilde{J}_N is proportional to

$$|\Delta(\tilde{\lambda})|^{2\kappa} \prod_{i=1}^N (\tilde{\lambda}_i^{\alpha} e^{-\tilde{\lambda}_i}), \quad \tilde{\lambda}_i > 0.$$

Let $\tilde{\mu}_N$ be the spectral measure of \tilde{J}_N and

$$m_r(N, \kappa, \alpha) = \mathbb{E}[\langle \tilde{\mu}_N, x^r \rangle] = \mathbb{E}[(\tilde{J}_N)^r(1, 1)].$$

Then $m_r(N, \kappa, \alpha)$ satisfies the following duality relation.

Lemma 3.4 (cf. [5, Theorem 2.11]). *The function $m_r(N, \kappa, \alpha)$ is a polynomial with respect to N, κ and α and satisfies the following relation*

$$m_r(N, \kappa, \alpha) = (-1)^r \kappa^r m_r(-\kappa N, \kappa^{-1}, -\alpha/\kappa).$$

Based on the above duality relation, we now identify the limit of $m_r(N, \kappa, \alpha)$ in the regime where $\kappa N \rightarrow c$. For fixed N , it is straightforward to calculate the limit of $\kappa^{-1/2} \tilde{B}_N$ with parameters $(N, \kappa, a\kappa)$ as $\kappa \rightarrow \infty$, where a is fixed,

$$\left(\frac{1}{\sqrt{\kappa}} \tilde{B}_N \right) \rightarrow \begin{pmatrix} \sqrt{a+N-1} & & & & \\ \sqrt{N-1} & \sqrt{a+N-2} & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \sqrt{1} & \sqrt{a} \end{pmatrix} =: D_N(a).$$

Here the convergence holds in L^q entry-wisely. Therefore,

$$\lim_{\kappa \rightarrow \infty} \kappa^{-r} m_r(N, \kappa, a\kappa) = \lim_{\kappa \rightarrow \infty} \mathbb{E}[\kappa^{-r} (\tilde{J}_N)^r(1, 1)] = (D_N(a) D_N(a)^t)^r(1, 1).$$

Next, in viewing of the duality relation, it holds that

$$\lim_{N \rightarrow \infty, \kappa N \rightarrow c} m_r(N, \kappa, \alpha) = \lim_{\kappa \rightarrow \infty} (-1)^r \kappa^{-r} m_r(-c, \kappa, -\alpha\kappa).$$

Let us consider the following infinite matrix by exchanging $N \leftrightarrow -c, a \leftrightarrow -\alpha$ and replacing the sign inside square roots as well,

$$W_{\alpha,c} = \begin{pmatrix} \sqrt{\alpha+c+1} & & & & \\ \sqrt{c+1} & \sqrt{\alpha+c+2} & & & \\ & \sqrt{c+2} & \sqrt{\alpha+c+3} & & \\ & & & \ddots & \\ & & & & \ddots \end{pmatrix}.$$

Let $J_{\alpha,c} = W_{\alpha,c} W_{\alpha,c}^t$ and $l_r(\alpha, c) = (J_{\alpha,c})^r(1, 1)$. Then it follows that

$$\lim_{\kappa N \rightarrow c} m_r(N, \kappa, \alpha) = \lim_{\kappa \rightarrow \infty} (-1)^r \kappa^{-r} m_r(-c, \kappa, -\alpha\kappa) = l_r(\alpha, c).$$

In conclusion, we have just proved the following result.

Theorem 3.5. *Let $\alpha > -1$ and $c \geq 0$ be given. Then in the regime where $\beta N \rightarrow 2c$,*

$$\mathbb{E} \left[\frac{1}{N} \text{tr}((\tilde{J}_N)^r) \right] = \mathbb{E}[(\tilde{J}_N)^r(1, 1)] \rightarrow \langle \mu_{\alpha,c}, x^r \rangle,$$

for any $r \in \mathbb{N}$. Here recall that $\mu_{\alpha,c}$ is the spectral measure of $J_{\alpha,c}$ whose density is given in Lemma 2.1

The limiting measure in this regime was calculated in [1] by a different method.

3.3. Almost sure convergence. In this section, we complete the proof of Theorem 1.1 by showing the following almost sure convergence.

Lemma 3.6. *For any $r \in \mathbb{N}$, as $N\beta \rightarrow 2c \in (0, \infty]$,*

$$S_N := \frac{1}{N} \operatorname{tr}[(J_N)^r] - \mathbb{E} \left[\frac{1}{N} \operatorname{tr}[(J_N)^r] \right] \rightarrow 0, \quad \text{almost surely.}$$

The idea is that for fixed r , $p_i := (J_N)^r(i, i)$ is independent of $p_j = (J_N)^r(j, j)$, if $|i - j| \geq D_r$, where D_r is a constant. Then write S_N as a sum of D_r sums of independent random variables

$$S_N = \frac{1}{N} \sum_i (p_{1+iD_r} - \mathbb{E}[p_{1+iD_r}]) + \cdots + \frac{1}{N} \sum_i (p_{D_r+iD_r} - \mathbb{E}[p_{D_r+iD_r}]).$$

For each sum of independent random variables, we use the following result whose proof can be found in the proof of Theorem 2.3 in [15] to show the almost sure convergence.

Lemma 3.7. *Assume that for each N , the random variables $\{\xi_{N,i}\}_{i=1}^{\ell_N}$ are independent and that*

$$(7) \quad \sup_N \sup_{1 \leq i \leq \ell_N} \mathbb{E}[(\xi_{N,i})^4] < \infty.$$

Assume further that $\ell_N/N \rightarrow \text{const} \in (0, \infty)$ as $N \rightarrow \infty$. Then as $N \rightarrow \infty$,

$$\frac{1}{N} \sum_{i=1}^{\ell_N} (\xi_{N,i} - \mathbb{E}[\xi_{N,i}]) \rightarrow 0, \quad \text{almost surely.}$$

The moment condition (7) can be easily checked for p_i in the regime $\beta N \rightarrow 2c \in (0, \infty]$. The desired almost sure convergence then follows immediately.

4. Gaussian fluctuations around the limit

4.1. Polynomial test functions. In this section, we establish central limit theorems (CLTs) for polynomial test functions. Since arguments are similar to those used in [15] in case of Gaussian beta ensembles, we only sketch main steps. Without loss of generality, assume that $M = N/\gamma$, where $\gamma \in (0, 1)$ is fixed.

Let us write B_N as

$$B_N = \frac{\sqrt{\gamma}}{\sqrt{\beta N}} \begin{pmatrix} c_1 & & & & & \\ d_1 & c_2 & & & & \\ & \ddots & \ddots & & & \\ & & & d_{N-1} & c_N & \\ & & & & & \end{pmatrix}, \quad \text{where } \begin{cases} c_i \sim \chi_{\frac{\beta N}{\gamma} - \beta(i-1)}, \\ d_i \sim \chi_{\beta N - \beta i}. \end{cases}$$

Then

$$J_N = B_N(B_N)^t = \frac{\gamma}{\beta N} \begin{pmatrix} c_1^2 & c_1 d_1 & & & & \\ c_1 d_1 & c_2^2 + d_1^2 & c_2 d_2 & & & \\ & \ddots & \ddots & \ddots & & \\ & & & c_{N-1} d_{N-1} & c_N^2 + d_{N-1}^2 & \\ & & & & & \end{pmatrix}.$$

Recall that for fixed $r \geq 1$, the r th moment $\langle \mu_N, x^r \rangle$ is a polynomial of $\{c_i, d_i\}_{1 \leq i \leq \frac{r+1}{2}}$. It is actually a polynomial of $\{c_i^2, d_i^2\}$, that is,

$$\langle \mu_N, x^r \rangle = (J_N)^r(1, 1) = \frac{\gamma^r}{(\beta N)^r} \sum_{\vec{\eta}, \vec{\zeta}} a(\vec{\eta}, \vec{\zeta}) \prod_{i=1}^r c_i^{2\eta_i} d_i^{2\zeta_i},$$

where non-negative integer vectors $\vec{\eta} = (\eta_1, \dots, \eta_r)$ and $\vec{\zeta} = (\zeta_1, \dots, \zeta_r)$ satisfy $\sum_{i=1}^r (\eta_i + \zeta_i) = r$. Then, from formulae for moments of the chi-squared distribution, we conclude that

Lemma 4.1. (i) For $r \in \mathbb{N}$,

$$\mathbb{E}[\langle \mu_N, x^r \rangle] = \sum_{k=0}^r \frac{R_{r,k}(\beta)}{(\beta N)^k},$$

where $R_{r,k}(\beta)$ is a polynomial in β of degree at most k .

(ii) For $r, s \in \mathbb{N}$,

$$\mathbb{E}[\langle \mu_N, x^r \rangle \langle \mu_N, x^s \rangle] = \sum_{k=0}^{r+s} \frac{Q_{r,s;k}(\beta)}{(\beta N)^k},$$

where $Q_{r,s;k}(\beta)$ is a polynomial in β of degree at most k .

Let p be a polynomial of degree m . From the above expressions, we can derive a general form of $\text{Var}[\langle \mu_N, p \rangle]$, and then that of $\text{Var}[\langle L_N, p \rangle]$ by taking into account of the relation (5). Similar to the case of Gaussian beta ensembles, the formula for $\text{Var}[\langle L_N, p \rangle]$ can be simplified as follows. The proof is omitted.

Lemma 4.2. Let p be a polynomial of order m . Then the variance $\text{Var}[\langle L_N, p \rangle]$ can be expressed as

$$\text{Var}[\langle L_N, p \rangle] = \sum_{k=2}^{2m+1} \frac{\beta \ell_{p;k}(\beta)}{(\beta N)^k},$$

where $\ell_{p;k}(\beta)$ is a polynomial in β of degree at most $(k - 2)$.

Corollary 4.3. (i) As $\beta N \rightarrow \infty$,

$$\beta N^2 \text{Var}[\langle L_N, p \rangle] \rightarrow \sigma_p^2.$$

(ii) As $\beta N \rightarrow 2c$,

$$\beta N^2 \text{Var}[\langle L_N, p \rangle] \rightarrow \sigma_{p,c}^2.$$

Theorem 3.4 in [15] provides sufficient conditions under which CLTs for $\{\langle L_N, p \rangle\}$ hold in case of Jacobi matrices with independent entries. The result can be easily extended to Wishart-type Jacobi matrices $\{J_N\}$ here by considering the filtration $\{\mathcal{F}_k = \sigma\{c_i, d_i : i = 1, \dots, k\}\}_k$. The convergence of variances as in the previous corollary is one of the two sufficient conditions. The remaining one is quite similar to that in the Gaussian beta ensembles case, and hence, we will not mention it in details here. Consequently, the following CLTs for polynomial test functions follow.

Theorem 4.4. Let p be a polynomial. Then the following hold.

(i) As $\beta N \rightarrow \infty$,

$$\sqrt{\beta N}(\langle L_N, p \rangle - \mathbb{E}[\langle L_N, p \rangle]) \xrightarrow{d} \mathcal{N}(0, \sigma_p^2).$$

(ii) As $\beta N \rightarrow 2c$,

$$\sqrt{\beta N}(\langle L_N, p \rangle - \mathbb{E}[\langle L_N, p \rangle]) \xrightarrow{d} \mathcal{N}(0, \sigma_{p,c}^2).$$

REMARK 4.5. (i) For fixed β , it was shown in [5] that as $N \rightarrow \infty$,

$$N\langle L_N, p \rangle - N\langle mp_\gamma, p \rangle - \left(\frac{2}{\beta} - 1\right)\langle \mu_1, p \rangle \xrightarrow{d} \mathcal{N}(0, \sigma_p^2/\beta),$$

where μ_1 is given by

$$\mu_1 = \frac{1}{4}\delta_{\lambda_-} + \frac{1}{4}\delta_{\lambda_+} + \frac{1}{2\pi} \frac{1}{\sqrt{(\lambda_+ - x)(x - \lambda_-)}} \mathbf{1}_{(\lambda_-, \lambda_+)}(dx).$$

(ii) Using results in case $\beta = 1, 2$, we deduce that the limiting variance in the regime $\beta N \rightarrow \infty$ is given by

$$\sigma_f^2 = \frac{1}{2\pi^2} \int_{\lambda_-}^{\lambda^+} \int_{\lambda_-}^{\lambda^+} \left(\frac{f(y) - f(x)}{y - x}\right)^2 \frac{4\gamma - (x - \lambda_m)(y - \lambda_m)}{\sqrt{4\gamma - (x - \lambda_m)^2} \sqrt{4\gamma - (y - \lambda_m)^2}} dx dy,$$

where $\lambda_m = (\lambda_- + \lambda_+)/2 = (1 + \gamma)$; cf. [12, Theorem 7.3.1].

4.2. C^1 test functions. To extend CLTs from polynomial test functions to C^1 test functions, one idea is to use a type of Poincaré inequality. Consider (scaled) β -Laguerre ensembles with the joint density proportional to

$$|\Delta(\lambda)|^\beta \prod_{i=1}^N (\lambda_i^\alpha e^{-\eta\lambda_i}),$$

where $\alpha = \frac{\beta}{2}(M - N + 1) - 1$, and $\eta > 0$. Directly from the joint density, we can derive a Poincaré inequality by using the following result. However, this approach requires $\alpha > 0$.

Lemma 4.6 ([3, Proposition 2.1]). *Let $dv = e^{-V} dx$ be a probability measure supported on an open convex set $\Omega \subset \mathbb{R}^N$. Assume that V is twice continuously differentiable and strictly convex on Ω . Then for any locally Lipschitz function F on Ω ,*

$$\text{Var}_v[F] = \int F^2 dv - \left(\int F dv\right)^2 \leq \int (\nabla F)^t \text{Hess}(V)^{-1} \nabla F dv.$$

Here $\text{Hess}(V)$ denotes the Hessian of V .

Let $\Omega = \{(\lambda_1, \dots, \lambda_N) : 0 < \lambda_1 < \dots < \lambda_N\} \subset \mathbb{R}^N$. Let ν be the distribution of the ordered eigenvalues of (scaled) β -Laguerre ensembles, that is, the probability measure on Ω of the form

$$d\nu = \text{const} \times |\Delta(\lambda)|^\beta \prod_{i=1}^N (\lambda_i^\alpha e^{-\eta\lambda_i}) d\lambda_1 \cdots d\lambda_N = e^{-V} d\lambda_1 \cdots d\lambda_N,$$

where

$$V = \text{const} + \eta \sum_{i=1}^N \lambda_i - \alpha \sum_{i=1}^N \log \lambda_i - \frac{\beta}{2} \sum_{i \neq j} \log |\lambda_j - \lambda_i|.$$

Then the Hessian matrix of V can be easily calculated

$$\frac{\partial^2 V}{\partial \lambda_i^2} = \frac{\alpha}{\lambda_i^2} + \beta \sum_{j \neq i} \frac{1}{(\lambda_j - \lambda_i)^2},$$

$$\frac{\partial^2 V}{\partial \lambda_i \partial \lambda_j} = -\beta \frac{1}{(\lambda_j - \lambda_i)^2}.$$

Observe that $\text{Hess}(V) \geq D = \text{diag}(\frac{\alpha}{\lambda_i^2})$. Here for two real symmetric matrices A and B , the notation $A \geq B$ indicates that $A - B$ is positive semi-definite. It follows that $\text{Hess}(V)^{-1} \leq D^{-1}$. And hence, using Lemma 4.6 with

$$F = \frac{1}{N} \sum_{i=1}^N f(\lambda_i),$$

for a continuously differentiable function f , we get the following inequality

$$\text{Var}_v[F] \leq \int (\nabla F)^t \text{Hess}(V)^{-1} \nabla F dv \leq \int (\nabla F)^t D^{-1} \nabla F dv.$$

The inequality can be rewritten as

$$(8) \quad \text{Var}[\langle L_N, f \rangle] \leq \frac{1}{\alpha N} \mathbb{E}[\langle L_N, (\lambda f'(\lambda))^2 \rangle].$$

This is one type of Poincaré inequality for β -Laguerre ensembles in this paper.

The restriction of the above inequality is that α must be positive, which does not hold in the regime $\beta N \rightarrow 2c$ when c is small. The second approach based on the random Jacobi matrix model removes such restriction. We will end up with a slightly different inequality. Let us begin with several concepts.

A real random variable X is said to satisfy a Poincaré inequality if there is a constant $c > 0$ such that for any smooth function $f: \mathbb{R} \rightarrow \mathbb{R}$,

$$\text{Var}[f(X)] \leq c \mathbb{E}[f'(X)^2].$$

Here, by smooth we mean enough regularity so that the considering terms make sense. By definition, it is clear that X satisfies a Poincaré inequality with a constant c , if and only if αX satisfies a Poincaré inequality with a constant $c\alpha^2$, for non-zero constant α .

Lemma 4.7. *For $k > 0$, the chi distribution χ_k satisfies a Poincaré inequality with $c = 1$, that is,*

$$\text{Var}[f(X)] \leq \mathbb{E}[f'(X)^2], \quad X \sim \chi_k.$$

Proof. The probability density function of the chi distribution with k degrees of freedom is given by

$$\frac{1}{2^{\frac{k}{2}-1} \Gamma(\frac{k}{2})} x^{k-1} e^{-\frac{x^2}{2}}, \quad (x > 0).$$

Thus, for $k \geq 1$, the conclusion follows immediately from Lemma 4.6.

Next, we consider the case $0 < k < 1$. Let $Y = X^k$. Then the probability density function of Y is given by

$$\text{const} \times \exp\left(-\frac{y^{\frac{2}{k}}}{2}\right), \quad (y > 0).$$

By using Lemma 4.6 with $V = const + \frac{y^{\frac{k}{2}}}{2}$, we obtain that

$$\text{Var}[g(Y)] \leq \frac{k^2}{2-k} \mathbb{E}[Y^{2-\frac{2}{k}} g'(Y)^2].$$

For given $f(x)$, let $g(y) = f(y^{\frac{1}{k}})$. Then we see that

$$\text{Var}[f(X)] = \text{Var}[g(Y)] \leq \frac{k^2}{2-k} \mathbb{E}[Y^{2-\frac{2}{k}} g'(Y)^2] = \frac{1}{2-k} \mathbb{E}[f'(X)^2] \leq \mathbb{E}[f'(X)^2].$$

This means that X satisfies a Poincaré inequality with a constant $c = 1$. The proof is complete. □

We need the following property.

Lemma 4.8 ([10, Corollary 5.7]). *Assume that $X_i, (i = 1, \dots, k)$, satisfy Poincaré inequalities with constants c_i . Assume further that $\{X_i\}$ are independent. Then for any smooth function $g: \mathbb{R}^k \rightarrow \mathbb{R}$,*

$$\text{Var}[g(X_1, \dots, X_k)] \leq (\max_i c_i) \mathbb{E}[|\nabla g(X_1, \dots, X_k)|^2].$$

Let $Y = (y_{mn})$ be an $M \times N$ real matrix, and $X = Y^t Y = (x_{ij})$. For a continuously differentiable function $f: \mathbb{R} \rightarrow \mathbb{R}$, let

$$g = g((y_{mn})) = \text{tr}(f(X)).$$

Then the partial derivatives of g can be expressed as follows.

Lemma 4.9 (cf. Eq. (7.2.5) in [12]). *It holds that*

$$\left(\frac{\partial g}{\partial y_{mn}} \right)_{M \times N} = 2Y f'(X).$$

Consequently

$$\sum_{m,n} \left(\frac{\partial g}{\partial y_{mn}} \right)^2 = 4 \text{tr}(Y f'(X) f'(X) Y^t) = 4 \text{tr}(X f'^2(X)) = 4 \sum_{i=1}^N \lambda_i f'(\lambda_i)^2.$$

Here $\lambda_1, \dots, \lambda_N$ are the eigenvalues of X .

Combining three lemmas above, we arrive at another type of Poincaré inequality for β -Laguerre ensembles.

Theorem 4.10. *Assume that f is a continuously differentiable function. Then for the β -Laguerre ensemble (1), the following inequality holds*

$$(9) \quad \text{Var}[\langle L_N, f \rangle] \leq \frac{4}{\beta MN} \mathbb{E}[\langle L_N, x f'(x)^2 \rangle].$$

The above Poincaré inequality is a key ingredient to extend CLTs to C^1 test functions. Let us now prove Theorem 1.2.

Proof of Theorem 1.2. Assume that f has continuous derivative f' of polynomial growth. This implies that $f' \in L^2((1 + x^2) d\nu_{\gamma,c}(x))$. Recall that for convenience, $\nu_{\gamma,\infty}$ denotes the Marchenko–Pastur distribution with parameter γ . We need the following property of mea-

asures determined by moments (called M. Riesz’s Theorem (1923) in [2]): the measure μ is determined by moments, if and only if polynomials are dense in $L^2((1 + x^2)d\mu(x))$. Consequently, there is a sequence of polynomials $\{p_k\}$ converging to f' in $L^2((1 + x^2)dv_{\gamma,c}(x))$. It then follows that

$$\int x(p_k - f')^2 dv_{\gamma,c}(x) \leq \frac{1}{2} \int (p_k - f')^2 (1 + x^2) dv_{\gamma,c}(x) \rightarrow 0 \quad \text{as } k \rightarrow \infty.$$

Let P_k be a primitive of p_k . Since P_k is a polynomial, for fixed k , as $N \rightarrow \infty$,

$$(10) \quad X_{N,k} := \sqrt{\beta N}(\langle L_N, P_k \rangle - \mathbb{E}[\langle L_N, P_k \rangle]) \xrightarrow{d} \mathcal{N}(0, \sigma_{P_k,c}^2), \quad \text{Var}[X_{N,k}] \rightarrow \sigma_{P_k,c}^2.$$

Let $Y_N = \sqrt{\beta N}(\langle L_N, f \rangle - \mathbb{E}[\langle L_N, f \rangle])$. Then by the Poincaré inequality (9),

$$(11) \quad \text{Var}[Y_N - X_{N,k}] \leq \frac{4N}{M} \mathbb{E}[\langle L_N, x(f' - p_k)^2 \rangle],$$

which implies that

$$\lim_{k \rightarrow \infty} \limsup_{N \rightarrow \infty} \text{Var}[Y_N - X_{N,k}] \leq 4\gamma \lim_{k \rightarrow \infty} \langle v_{\gamma,c}, x(f' - p_k)^2 \rangle = 0.$$

Here we have used the property that for continuous function g of polynomial growth,

$$\mathbb{E}[\langle L_N, g \rangle] \rightarrow \langle v_{\gamma,c}, g \rangle.$$

Then, the CLT for Y_N follows from the equations (10) and (11) by using the following general result whose proof can be found in [11] or [16]. The proof is complete. \square

Lemma 4.11. *Let $\{Y_N\}_{N=1}^\infty$ and $\{X_{N,k}\}_{N,k=1}^\infty$ be mean zero real random variables. Assume that*

- (i) *for each k , as $N \rightarrow \infty$, $X_{N,k} \xrightarrow{d} \mathcal{N}(0, \sigma_k^2)$, and $\text{Var}[X_{N,k}] \rightarrow \sigma_k^2$;*
- (ii) *$\lim_{k \rightarrow \infty} \limsup_{N \rightarrow \infty} \text{Var}[X_{N,k} - Y_N] = 0$.*

Then the limit $\sigma^2 = \lim_{k \rightarrow \infty} \sigma_k^2$ exists, and as $N \rightarrow \infty$,

$$Y_N \xrightarrow{d} \mathcal{N}(0, \sigma^2), \quad \text{Var}[Y_N] \rightarrow \sigma^2.$$

ACKNOWLEDGEMENTS. The authors would like to thank the referees for helpful comments. This work is supported by University of Science, Vietnam National University, Hanoi under project number TN.18.03 (H.D.T) and by JSPS KAKENHI Grant Number JP19K14547 (K.D.T.).

References

- [1] R. Allez, J.P. Bouchaud, S.N. Majumdar and P. Vivo: *Invariant β -Wishart ensembles, crossover densities and asymptotic corrections to the Marnenko-Pastur law*, J. Phys. A **46** (2013), 015001, 22pp.
- [2] A.G. Bakan: *Polynomial density in $L_p(\mathbb{R}, d\mu)$ and representation of all measures which generate a determinate Hamburger moment problem*; in Approximation, optimization and mathematical economics (Pointe-à-Pitre, 1999), Physica, Heidelberg, 2001, 37–46.

- [3] S.G. Bobkov and M. Ledoux: *From Brunn-Minkowski to Brascamp-Lieb and to logarithmic Sobolev inequalities*, *Geom. Funct. Anal.* **10** (2000), 1028–1052.
- [4] I. Dumitriu and A. Edelman: *Matrix models for beta ensembles*, *J. Math. Phys.* **43** (2002), 5830–5847.
- [5] I. Dumitriu and A. Edelman: *Global spectrum fluctuations for the β -Hermite and β -Laguerre ensembles via matrix models*, *J. Math. Phys.* **47** (2006), 063302, 36pp.
- [6] T.K. Duy: *On spectral measures of random Jacobi matrices*, *Osaka J. Math.* **55** (2018), 595–617.
- [7] T.K. Duy and T. Shirai: *The mean spectral measures of random Jacobi matrices related to Gaussian beta ensembles*, *Electron. Commun. Probab.* **20** (2015), no. 68, 13pp.
- [8] M.E.H. Ismail, J. Letessier and G. Valent: *Linear birth and death models and associated Laguerre and Meixner polynomials*, *J. Approx. Theory* **55** (1988), 337–348.
- [9] S. Jacquot and B. Valkó: *Bulk scaling limit of the Laguerre ensemble*, *Electron. J. Probab.* **16** (2011), 314–346.
- [10] M. Ledoux: *The Concentration of Measure Phenomenon*, *Mathematical Surveys and Monographs* **89**, American Mathematical Society, Providence, RI, 2001.
- [11] F. Nakano and K.D. Trinh: *Gaussian beta ensembles at high temperature: eigenvalue fluctuations and bulk statistics*, *J. Stat. Phys.* **173** (2018), 295–321.
- [12] L. Pastur and M. Shcherbina: *Eigenvalue Distribution of Large Random Matrices*, *Mathematical Surveys and Monographs* **171**, American Mathematical Society, Providence, RI, 2011.
- [13] B. Simon: *Szegő's Theorem and its Descendants: Spectral Theory for L^2 Perturbations of Orthogonal Polynomials*, *M.B. Porter Lectures*, Princeton University Press, Princeton, NJ, 2011.
- [14] P. Sosoe and P. Wong: *Convergence of the eigenvalue density for β -Laguerre ensembles on short scales*, *Electron. J. Probab.* **19** (2014), no. 34, 18pp.
- [15] K.D. Trinh: *Global spectrum fluctuations for Gaussian beta ensembles: a martingale approach*, *J. Theoret. Probab.* **32** (2019), 1420–1437.
- [16] K.D. Trinh: *On central limit theorems in stochastic geometry for add-one cost stabilizing functionals*, *Electron. Commun. Probab.* **24** (2019), no. 76, 15 pp.

Hoang Dung Trinh
 Faculty of Mathematics Mechanics Informatics
 University of Science
 Vietnam National University
 Hanoi
 Vietnam
 e-mail: thdung.hus@gmail.com

Khanh Duy Trinh
 Global Center for Science and Engineering
 Waseda University
 Tokyo
 Japan
 e-mail: trinh@aoni.waseda.jp